

Datasheet for ABIN651929
anti-RNF208 antibody (AA 86-115)[Go to Product page](#)

3 Images

Overview

Quantity:	400 µL
Target:	RNF208
Binding Specificity:	AA 86-115
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This RNF208 antibody is un-conjugated
Application:	Western Blotting (WB), Flow Cytometry (FACS), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

Product Details

Immunogen:	This RNF208 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 86-115 amino acids from the Central region of human RNF208.
Clone:	RB27242
Isotype:	Ig Fraction
Purification:	This antibody is purified through a protein A column, followed by peptide affinity purification.

Target Details

Target:	RNF208
Alternative Name:	RNF208 (RNF208 Products)

Target Details

Molecular Weight:	27964
Gene ID:	727800
NCBI Accession:	NP_112587
UniProt:	Q9H0X6


Application Details

Application Notes:	WB: 1:1000. IHC-P: 1:50~100. FC: 1:10~50
Restrictions:	For Research Use only

Handling


Format:	Liquid
Buffer:	Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Maintain refrigerated at 2-8 °C for up to 6 months. For long term storage store at -20 °C in small aliquots to prevent freeze-thaw cycles.
Expiry Date:	6 months

Images



Flow Cytometry

Image 1. RN Antibody (Center) (ABIN651929 and ABIN2840458) flow cytometric analysis of HepG2 cells (bottom histogram) compared to a negative control cell (top histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.



Western Blotting

Image 2. Western blot analysis of lysate from A431 cell line, using RN Antibody (Center) (ABIN651929 and ABIN2840458). (ABIN651929 and ABIN2840458) was diluted at 1:1000 at each lane. A goat anti-rabbit IgG H&L(HRP) at 1:5000 dilution was used as the secondary antibody. Lysate at 35 µg per lane.

Immunohistochemistry (Paraffin-embedded Sections)

Image 3. RN Antibody (Center) (ABIN651929 and ABIN2840458) immunohistochemistry analysis in formalin fixed and paraffin embedded human brain tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the RN Antibody (Center) for immunohistochemistry. Clinical relevance has not been evaluated.