

Datasheet for ABIN652043

anti-ADCY10 antibody (AA 557-583)**2** Images**1** Publication[Go to Product page](#)

Overview

Quantity:	400 µL
Target:	ADCY10
Binding Specificity:	AA 557-583
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This ADCY10 antibody is un-conjugated
Application:	Western Blotting (WB), Flow Cytometry (FACS)

Product Details

Immunogen:	This ADCY10 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 557-583 amino acids from the Central region of human ADCY10.
Clone:	RB22128
Isotype:	Ig Fraction
Purification:	This antibody is purified through a protein A column, followed by peptide affinity purification.

Target Details

Target:	ADCY10
Alternative Name:	ADCY10 (ADCY10 Products)
Molecular Weight:	187149

Target Details

Gene ID:	55811
NCBI Accession:	NP_001161221 , NP_060887
UniProt:	Q96PN6
Pathways:	cAMP Metabolic Process

Application Details


Application Notes:	WB: 1:1000. FC: 1:10~50
Restrictions:	For Research Use only

Handling

Format:	Liquid
Buffer:	Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C, -20 °C
Storage Comment:	Maintain refrigerated at 2-8 °C for up to 6 months. For long term storage store at -20 °C in small aliquots to prevent freeze-thaw cycles.
Expiry Date:	6 months

Publications

Product cited in:	Chen, Chen, Deng, Hao, Wang, Liu: "Sodium-hydrogen exchanger NHA1 and NHA2 control sperm motility and male fertility." in: Cell death & disease , Vol. 7, pp. e2152, (2016) (PubMed).
-------------------	--



Flow Cytometry

Image 1. ADCY10 Antibody (Center) (ABIN652043 and ABIN2840517) flow cytometric analysis of K562 cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

Western Blotting

Image 2. Anti-ADCY10 Antibody (Center) at 1:1000 dilution + K562 whole cell lysate Lysates/proteins at 20 μ g per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 187 kDa Blocking/Dilution buffer: 5 % NFDm/TBST.