# antibodies -online.com









Go to Product page

$\sim$					
	1//6	r	<b>V</b> I	$\Theta$	Λ

Quantity:	400 μL
Target:	DIRAS3
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This DIRAS3 antibody is un-conjugated
Application:	Western Blotting (WB), Flow Cytometry (FACS), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

### **Product Details**

Immunogen:	This DIRAS3 Antibody is generated from rabbits immunized with recombinant protein of human DIRAS3.
Clone:	RB23150
Isotype:	lg Fraction
Purification:	This antibody is purified through a protein A column, followed by peptide affinity purification.
Target Details	

### Larget Details

Target:	DIRAS3
Alternative Name:	DIRAS3 (DIRAS3 Products)
Molecular Weight:	25861

# **Target Details**

Gene ID:	9077
NCBI Accession:	NP_004666
UniProt:	O95661


# **Application Details**

Application Notes:	WB: 1:1000. WB: 1:1000. IHC-P: 1:50~100. FC: 1:10~50
Restrictions:	For Research Use only

# Handling


Format:	Liquid
Buffer:	Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Maintain refrigerated at 2-8 °C for up to 6 months. For long term storage store at -20 °C in small aliquots to prevent freeze-thaw cycles.
Expiry Date:	6 months


## **Images**



# **Flow Cytometry**

**Image 1.** DIRAS3 Antibody (ABIN652072 and ABIN2840536) flow cytometric analysis of WiDr cells (right histogram) compared to a negative control cell (left histogram).FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.





### **Western Blotting**

Image 2. Western blot analysis of lysates from Hela cell line and human ovary tissue lysate (from left to right), using DIRAS3 Antibody (ABIN652072 and ABIN2840536). (ABIN652072 and ABIN2840536) was diluted at 1:1000 at each lane. A goat anti-rabbit IgG H&L(HRP) at 1:10000 dilution was used as the secondary antibody. Lysates at 35 μg per lane.

### **Western Blotting**

**Image 3.** DIRAS3 Antibody (ABIN652072 and ABIN2840536) western blot analysis in WiDr cell line lysates (35  $\mu$  g/lane). This demonstrates the DIRAS3 antibody detected the DIRAS3 protein (arrow).

Please check the product details page for more images. Overall 4 images are available for ABIN652072.