antibodies -online.com





Datasheet for ABIN652176

anti-TRAM2 antibody (C-Term)

2 Images



Go to Product page

Overview

Overview	
Quantity:	400 μL
Target:	TRAM2
Binding Specificity:	AA 342-370, C-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This TRAM2 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))
Product Details	
Immunogen:	This TRAM2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 342-370 amino acids from the C-terminal region of human TRAM2.
Clone:	RB27529
Isotype:	lg Fraction
Purification:	This antibody is purified through a protein A column, followed by peptide affinity purification.
Target Details	
Target:	TRAM2
Alternative Name:	TRAM2 (TRAM2 Products)
Molecular Weight:	43328

Target Details

Gene ID:	9697
NCBI Accession:	NP_036420
UniProt:	Q15035


Application Details

Application Notes:	WB: 1:1000. IHC-P: 1:50~100
Restrictions:	For Research Use only

Handling


Format:	Liquid
Buffer:	Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Maintain refrigerated at 2-8 °C for up to 6 months. For long term storage store at -20 °C in small aliquots to prevent freeze-thaw cycles.
Expiry Date:	6 months

Images



Immunohistochemistry (Paraffin-embedded Sections)

Image 1. TR antibody (C-term) (ABIN652176 and ABIN2840578) immunohistochemistry analysis in formalin fixed and paraffin embedded human prostate carcinoma followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the TR antibody (C-term) for immunohistochemistry. Clinical relevance has not been evaluated.



Western Blotting

Image 2. TR Antibody (C-term) (ABIN652176 and ABIN2840578) western blot analysis in MDA-M cell line lysates (35 μ g/lane).This demonstrates the TR antibody detected the TR protein (arrow).