

Datasheet for ABIN652283
anti-AKT1 antibody (C-Term)[Go to Product page](#)

4 Images

3 Publications

Overview

Quantity:	400 µL
Target:	AKT1
Binding Specificity:	AA 438-468, C-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This AKT1 antibody is un-conjugated
Application:	Western Blotting (WB), Flow Cytometry (FACS)

Product Details

Immunogen:	This AKT1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 438-468 amino acids from the C-terminal region of human AKT1.
Clone:	RB3023-3024
Isotype:	Ig Fraction
Predicted Reactivity:	M, Rat
Purification:	This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Target Details

Target:	AKT1
---------	------

Target Details

Alternative Name:	AKT1 (AKT1 Products)
Background:	The serine-threonine protein kinase encoded by the AKT1 gene is catalytically inactive in serum-starved primary and immortalized fibroblasts. AKT1 and the related AKT2 are activated by platelet-derived growth factor. The activation is rapid and specific, and it is abrogated by mutations in the pleckstrin homology domain of AKT1. It was shown that the activation occurs through phosphatidylinositol 3-kinase. In the developing nervous system AKT is a critical mediator of growth factor-induced neuronal survival. Survival factors can suppress apoptosis in a transcription-independent manner by activating the serine/threonine kinase AKT1, which then phosphorylates and inactivates components of the apoptotic machinery.
Molecular Weight:	55686
Gene ID:	207
NCBI Accession:	NP_001014431 , NP_001014432 , NP_005154
UniProt:	P31749
Pathways:	PI3K-Akt Signaling , RTK Signaling , TCR Signaling , AMPK Signaling , Interferon-gamma Pathway , TLR Signaling , Fc-epsilon Receptor Signaling Pathway , EGFR Signaling Pathway , Neurotrophin Signaling Pathway , Response to Water Deprivation , Regulation of Actin Filament Polymerization , Carbohydrate Homeostasis , Glycosaminoglycan Metabolic Process , Cellular Glucan Metabolic Process , Regulation of Muscle Cell Differentiation , Cell-Cell Junction Organization , Regulation of Cell Size , Skeletal Muscle Fiber Development , Regulation of Carbohydrate Metabolic Process , Hepatitis C , Protein targeting to Nucleus , CXCR4-mediated Signaling Events , Signaling Events mediated by VEGFR1 and VEGFR2 , Negative Regulation of intrinsic apoptotic Signaling , Thromboxane A2 Receptor Signaling , Signaling of Hepatocyte Growth Factor Receptor , Positive Regulation of fat Cell Differentiation , VEGFR1 Specific Signals , VEGF Signaling , Warburg Effect

Application Details

Application Notes:	WB: 1:1000. WB: 1:1000. WB: 1:1000. FC: 1:10~50
Restrictions:	For Research Use only

Handling

Format:	Liquid
Buffer:	Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.
Preservative:	Sodium azide


Handling

Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Maintain refrigerated at 2-8 °C for up to 6 months. For long term storage store at -20 °C in small aliquots to prevent freeze-thaw cycles.
Expiry Date:	6 months

Publications


Product cited in:	Smith, Wietgreffe, Shang, Reilly, Southern, Perkey, Duan, Kohler, Müller, Robinson, Carlis, Li, Johnson, Haase: "Live simian immunodeficiency virus vaccine correlate of protection: immune complex-inhibitory Fc receptor interactions that reduce target cell availability." in: Journal of immunology (Baltimore, Md. : 1950) , Vol. 193, Issue 6, pp. 3126-33, (2014) (PubMed).
-------------------	--

Images




Flow Cytometry

Image 1. AKT1 Antibody (C-term) (ABIN652283 and ABIN2841190) flow cytometric analysis of MDA-M cells (right histogram) compared to a negative control cell (left histogram).FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.



Western Blotting

Image 2. Western blot analysis of hAKT1- (ABIN652283 and ABIN2841190) in Y79 cell line lysates (35 µg/lane). AKT1 (arrow) was detected using the purified Pab.



Western Blotting

Image 3. Western blot analysis of AKT1 (arrow) using rabbit polyclonal hAKT1- (ABIN652283 and ABIN2841190). 293 cell lysates (2 µg/lane) either nontransfected (Lane 1) or transiently transfected with the AKT1 gene (Lane 2) (Origene Technologies).

Please check the [product details page](#) for more images. Overall 4 images are available for ABIN652283.