

Datasheet for ABIN652376
anti-KIT antibody (AA 680-711)



[Go to Product page](#)

2 Images

Overview

Quantity:	400 µL
Target:	KIT
Binding Specificity:	AA 680-711
Reactivity:	Human, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This KIT antibody is un-conjugated
Application:	Western Blotting (WB), Flow Cytometry (FACS)

Product Details

Immunogen:	This KIT antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 680-711 amino acids from human KIT.
Clone:	RB15541
Isotype:	Ig Fraction
Predicted Reactivity:	Pig
Purification:	This antibody is purified through a protein A column, followed by peptide affinity purification.

Target Details

Target:	KIT
Alternative Name:	KIT (KIT Products)

Target Details


Background:	KIT is the human homolog of the proto-oncogene c-kit. C-kit was first identified as the cellular homolog of the feline sarcoma viral oncogene v-kit. KIT is a type 3 transmembrane receptor for MGF (mast cell growth factor, also known as stem cell factor).
Molecular Weight:	109865
Gene ID:	3815
NCBI Accession:	NP_000213 , NP_001087241
UniProt:	P10721
Pathways:	RTK Signaling , Fc-epsilon Receptor Signaling Pathway , EGFR Signaling Pathway , Neurotrophin Signaling Pathway , Sensory Perception of Sound , Stem Cell Maintenance , Production of Molecular Mediator of Immune Response , Regulation of long-term Neuronal Synaptic Plasticity

Application Details

Application Notes:	WB: 1:1000. FC: 1:10~50
Restrictions:	For Research Use only

Handling

Format:	Liquid
Buffer:	Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C, -20 °C
Storage Comment:	Maintain refrigerated at 2-8 °C for up to 6 months. For long term storage store at -20 °C in small aliquots to prevent freeze-thaw cycles.
Expiry Date:	6 months



Flow Cytometry

Image 1. KIT Antibody g flow cytometric analysis of 293 cells (right histogram) compared to a negative control cell (left histogram).FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

Western Blotting

Image 2. Western blot analysis of KIT Antibody g in mouse cerebellum tissue lysates (35 µg/lane). KIT (arrow) was detected using the purified Pab.