antibodies -online.com





anti-CD94 antibody (N-Term)

3 Images



Go to Product page

\sim				
	11/6	٦r١	/10	۱۸.

Quantity:	400 μL
Target:	CD94 (KLRD1)
Binding Specificity:	AA 31-57, N-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CD94 antibody is un-conjugated
Application:	Western Blotting (WB), Flow Cytometry (FACS), Immunohistochemistry (Paraffin-embedded
	Sections) (IHC (p))
Product Details	
Immunogen:	This KLRD1 antibody is generated from rabbits immunized with a KLH conjugated synthetic
	peptide between 31-57 amino acids from the N-terminal region of human KLRD1.
Clone:	RB21999
Isotype:	lg Fraction
Purification:	This antibody is purified through a protein A column, followed by peptide affinity purification.
Target Details	
Target:	CD94 (KLRD1)
Alternative Name:	KLRD1 (KLRD1 Products)

Target Details


Background:	KLRD1 plays a role as a receptor for the recognition of MHC class I HLA-E molecules by NK cells and some cytotoxic T-cells.
Molecular Weight:	20513
Gene ID:	3824
NCBI Accession:	NP_001107868, NP_002253, NP_031360
UniProt:	Q13241

Application Details

Application Notes:	WB: 1:1000. IHC-P: 1:50~100. FC: 1:10~50
Restrictions:	For Research Use only


Handling

Format:	Liquid
Buffer:	Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Maintain refrigerated at 2-8 °C for up to 6 months. For long term storage store at -20 °C in small aliquots to prevent freeze-thaw cycles.
Expiry Date:	6 months




Flow Cytometry

Image 1. Flow cytometric analysis of CEM cells using KLRD1 Antibody (N-term)(bottom histogram) compared to a negative control cell (top histogram). FITC-conjugated goatanti-rabbit secondary antibodies were used for the analysis.



Western Blotting

Image 2. Western blot analysis of KLRD1 Antibody (N-term) (ABIN652571 and ABIN2842384) in K562 cell line lysates (35 μ g/lane). KLRD1 (arrow) was detected using the purified Pab.



Immunohistochemistry (Paraffin-embedded Sections)

Image 3. Formalin-fixed and paraffin-embedded human lymph with KLRD1 Antibody (N-term), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry, clinical relevance has not been evaluated.