# antibodies -online.com





## anti-PPID antibody (N-Term)





Go to Product page

$\sim$						
	1//	Д	r۱	/1	$\triangle$	٨

400 μL
PPID
AA 85-115, N-Term
Human, Mouse
Rabbit
Polyclonal
This PPID antibody is un-conjugated
Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Flow
Cytometry (FACS)
This PPID antibody is generated from rabbits immunized with a KLH conjugated synthetic
peptide between 85-115 amino acids from the N-terminal region of human PPID.
RB17211
lg Fraction
Rat
This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by
dialysis against PBS.
PPID

### **Target Details**


Alternative Name:	PPID (PPID Products)
Background:	PPlases accelerate the folding of proteins. It catalyzes the cis-trans isomerization of proline imidic peptide bonds in oligopeptides.
Molecular Weight:	40764
Gene ID:	5481
NCBI Accession:	NP_005029
UniProt:	Q08752
Pathways:	Nuclear Hormone Receptor Binding


## Application Details


Application Notes:	WB: 1:1000. WB: 1:1000. IHC-P: 1:50~100. FC: 1:10~50
Restrictions:	For Research Use only

## Handling

Format:	Liquid	
Buffer:	Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.	
Preservative:	Sodium azide	
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.	
Storage:	4 °C,-20 °C	
Storage Comment:	Maintain refrigerated at 2-8 °C for up to 6 months. For long term storage store at -20 °C in small aliquots to prevent freeze-thaw cycles.	
Expiry Date:	6 months	







#### **Western Blotting**

**Image 1.** Western blot analysis of ID (arrow) using rabbit polyclonal ID Antibody (N-term) (ABIN652735 and ABIN2842487). 293 cell lysates (2 μg/lane) either nontransfected (Lane 1) or transiently transfected (Lane 2) with the ID gene.

#### **Western Blotting**

**Image 2.** Western blot analysis of ID Antibody (N-term) (ABIN652735 and ABIN2842487) in NIH-3T3 cell line lysates (35 µg/lane). ID (arrow) was detected using the purified Pab.

#### **Immunohistochemistry (Paraffin-embedded Sections)**

**Image 3.** Formalin-fixed and paraffin-embedded human brain tissue reacted with ID Antibody (N-term), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry, clinical relevance has not been evaluated.

Please check the product details page for more images. Overall 4 images are available for ABIN652735.