

Datasheet for ABIN653223
anti-TSPEAR antibody (AA 293-320)

3 Images

1 Publication

[Go to Product page](#)

Overview

Quantity:	400 µL
Target:	TSPEAR
Binding Specificity:	AA 293-320
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This TSPEAR antibody is un-conjugated
Application:	Western Blotting (WB), Flow Cytometry (FACS), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

Product Details

Immunogen:	This C21orf29 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 293-320 amino acids from the Central region of human C21orf29.
Clone:	RB23103
Isotype:	Ig Fraction
Purification:	This antibody is purified through a protein A column, followed by peptide affinity purification.

Target Details

Target:	TSPEAR
Alternative Name:	C21orf29 (TSPEAR Products)

Target Details

Molecular Weight:	74924
Gene ID:	54084
NCBI Accession:	NP_001258966 , NP_659428
UniProt:	Q8WU66
Pathways:	Sensory Perception of Sound

Application Details




Application Notes:	WB: 1:1000. IHC-P: 1:50~100. FC: 1:10~50
Restrictions:	For Research Use only

Handling

Format:	Liquid
Buffer:	Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Maintain refrigerated at 2-8 °C for up to 6 months. For long term storage store at -20 °C in small aliquots to prevent freeze-thaw cycles.
Expiry Date:	6 months

Publications

Product cited in:	Wei, Cheng, Ouyang, Zhang, Hu, Chen, Wang, Lu, Zhang, Wang, Liu: "Glycoprotein screening in colorectal cancer based on differentially expressed Tn antigen." in: Oncology reports , Vol. 36, Issue 3, pp. 1313-24, (2017) (PubMed).
-------------------	--



Flow Cytometry

Image 1. C21orf29 Antibody (Center) (ABIN653223 and ABIN2842758) flow cytometric analysis of K562 cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

Western Blotting

Image 2. Anti-C21orf29 Antibody (Center) at 1:1000 dilution + 293 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 75 kDa. Blocking/Dilution buffer: 5 % NFDM/TBST.

Immunohistochemistry (Paraffin-embedded Sections)

Image 3. Formalin-fixed and paraffin-embedded human Lung carcinoma reacted with C21orf29 Antibody (Center), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry, clinical relevance has not been evaluated.