

Datasheet for ABIN653260

anti-FLT1 antibody





Go to Product page

()	11/0	K / /	$\triangle V$

Quantity:	400 μL	
Target:	FLT1	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This FLT1 antibody is un-conjugated	
Application:	Western Blotting (WB), Flow Cytometry (FACS), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))	

Product Details

lmmunogen:	This VGFR1 antibody is generated from rabbits immunized with human VGFR1 recombinant protein.
Clone:	RB16736
Isotype:	lg Fraction
Purification:	This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Target Details

Target:	FLT1	
Alternative Name:	VGFR1 (FLT1 Products)	
Background:	round: VGFR1 encodes a member of the vascular endothelial growth factor receptor (VEGFR) family	

t,
? -

	trophoblast cells and peripheral blood monocytes.	
Molecular Weight:	150769	
Gene ID:	2321	
NCBI Accession:	NP_001153392, NP_001153502, NP_001153503, NP_002010	
UniProt:	P17948	
Pathways:	RTK Signaling, Signaling Events mediated by VEGFR1 and VEGFR2, VEGFR1 Specific Signals	

Application Details

Application Notes:	WB: 1:1000. IHC-P: 1:50~100. FC: 1:10~50
Restrictions:	For Research Use only
Handling	

Format:	Liquid
Buffer:	Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Maintain refrigerated at 2-8 °C for up to 6 months. For long term storage store at -20 °C in small aliquots to prevent freeze-thaw cycles.
Expiry Date:	6 months


Publications

Product cited in:

Dang, Zhang, Li, Hu, Li, Li, Wei, Shi: "High VEGFR1/2 expression levels are predictors of poor survival in patients with cervical cancer." in: **Medicine**, Vol. 96, Issue 1, pp. e5772, (2017) (


PubMed).

Images




Flow Cytometry

Image 1. VGFR1 Antibody (ABIN653260 and ABIN2842780) flow cytometric analysis of MDA-M cells (bottom histogram) compared to a negative control cell (top histogram)FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.



Western Blotting

Image 2. Western blot analysis of VGFR1 Antibody (ABIN653260 and ABIN2842780) in MCF-7 cell line lysates (35 μ g/lane). VGFR1 (arrow) was detected using the purified Pab.



Immunohistochemistry (Paraffin-embedded Sections)

Image 3. Formalin-fixed and paraffin-embedded human breast carcinoma reacted with VGFR1 Antibody, which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry, clinical relevance has not been evaluated.