antibodies - online.com







anti-MORG1 antibody (AA 126-154)



Image



\sim			
	N/6	1//r	$I \cap V$

Overview		
Quantity:	400 μL	
Target:	MORG1 (wdr83)	
Binding Specificity:	AA 126-154	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This MORG1 antibody is un-conjugated	
Application:	Western Blotting (WB)	
Product Details		
Immunogen:	This MORG1 antibody is generated from rabbits immunized with a KLH conjugated synthetic	
	peptide between 126-154 amino acids from the Central region of human MORG1.	
Clone:	RB24289	
Isotype:	Ig Fraction	
Purification:	This antibody is purified through a protein A column, followed by peptide affinity purification.	
Target Details		
Target:	MORG1 (wdr83)	
Alternative Name:	MORG1 (wdr83 Products)	
Background:	MORG1 is a member of the WD-40 protein family. The protein is proposed to function as a	

Target Details

molecular scaffold for various multimeric protein complexes. The protein associates with several components of the extracellular signal-regulated kinase (ERK) pathway, and promotes ERK activity in response to serum or other signals. The protein also interacts with egl nine homolog 3 (EGLN3, also known as PHD3) and regulates expression of hypoxia-inducible factor 1, and has been purified as part of the spliceosome.

Molecular Weight: 34343

Gene ID: 84292

NCBI Accession: NP_001093207, NP_115708

UniProt: Q9BRX9

Application Details

Application Notes: WB: 1:1000

Restrictions: For Research Use only

Handling

Format:

Buffer:

Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.

Preservative:

Sodium azide

This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.


Storage:

4 °C,-20 °C

Maintain refrigerated at 2-8 °C for up to 6 months. For long term storage store at -20 °C in small

aliquots to prevent freeze-thaw cycles.

Expiry Date: 6 months



Western Blotting

Image 1. Western blot analysis of MORG1 Antibody (Center) (ABIN653527 and ABIN2842925) in K562 cell line lysates (35 μ g/lane). MORG1 (arrow) was detected using the purified Pab.