.-online.com antibodies

Datasheet for ABIN653665 anti-GDA antibody (N-Term)

2 Images



Overview

Quantity:	400 µL
Target:	GDA
Binding Specificity:	AA 94-122, N-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This GDA antibody is un-conjugated
Application:	Western Blotting (WB), Flow Cytometry (FACS)

Product Details

Immunogen:	This GDA antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 94-122 amino acids from the N-terminal region of human GDA.
Clone:	RB23974
Isotype:	Ig Fraction
Purification:	This antibody is purified through a protein A column, followed by peptide affinity purification.


Target Details

Target:	GDA
Alternative Name:	GDA (GDA Products)
Background:	GDA is an enzyme that catalyzes the hydrolytic deamination of guanine, producing xanthine and

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/3 | Product datasheet for ABIN653665 | 09/12/2023 | Copyright antibodies-online. All rights reserved.

Target Details

	ammonia.
Molecular Weight:	51003
Gene ID:	9615
NCBI Accession:	NP_001229434, NP_001229435, NP_001229436, NP_004284
UniProt:	Q9Y2T3
Application Details	
Application Notes:	WB: 1:1000. FC: 1:10~50
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which
	should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Maintain refrigerated at 2-8 °C for up to 6 months. For long term storage store at -20 °C in small
	aliquots to prevent freeze-thaw cycles.
Expiry Date:	6 months



Flow Cytometry

Image 1. GDA Antibody (N-term) (ABIN653665 and ABIN2842999) flow cytometric analysis of HL-60 cells (right histogram) compared to a negative control cell (left histogram).FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

Western Blotting

Image 2. Western blot analysis of GDA Antibody (N-term) (ABIN653665 and ABIN2842999) in HL-60 cell line lysates (35 µg/lane). GDA (arrow) was detected using the purified Pab.

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 3/3 | Product datasheet for ABIN653665 | 09/12/2023 | Copyright antibodies-online. All rights reserved.