



[Go to Product page](#)

Datasheet for ABIN653686

anti-MAPK1IP1L antibody (AA 108-137)

2 Images

Overview

Quantity:	400 µL
Target:	MAPK1IP1L
Binding Specificity:	AA 108-137
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This MAPK1IP1L antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

Product Details

Immunogen:	This MAPK1IP1L antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 108-137 amino acids from the Central region of human MAPK1IP1L.
Clone:	RB24363
Isotype:	Ig Fraction
Predicted Reactivity:	B, Pr
Purification:	This antibody is purified through a protein A column, followed by peptide affinity purification.

Target Details

Target:	MAPK1IP1L
Alternative Name:	MAPK1IP1L (MAPK1IP1L Products)

Target Details

Molecular Weight:	24269
Gene ID:	93487
NCBI Accession:	NP_653179
UniProt:	Q8NDC0


Application Details

Application Notes:	WB: 1:1000. IHC-P: 1:50~100
Restrictions:	For Research Use only

Handling


Format:	Liquid
Buffer:	Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Maintain refrigerated at 2-8 °C for up to 6 months. For long term storage store at -20 °C in small aliquots to prevent freeze-thaw cycles.
Expiry Date:	6 months

Images



Immunohistochemistry (Paraffin-embedded Sections)

Image 1. PK1IP1L Antibody (Center) (ABIN653686 and ABIN2843011) IHC analysis in forlin fixed and paraffin embedded kidney carcino followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the PK1IP1L Antibody (Center) for immunohistochemistry. Clinical relevance has not been evaluated.



Western Blotting

Image 2. All lanes : Anti-PK1IP1L Antibody (Center) at 1:1000 dilution Lane 1: U-251 MG whole cell lysate Lane 2: Y79 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 24 kDa Blocking/Dilution buffer: 5 % NFDN/TBST.