antibodies -online.com







anti-ACSBG2 antibody (AA 237-265)





\sim					
	1//	r۱	/1	\triangle	٨

Overview	
Quantity:	400 μL
Target:	ACSBG2
Binding Specificity:	AA 237-265
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This ACSBG2 antibody is un-conjugated
Application:	Western Blotting (WB), Flow Cytometry (FACS), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))
Product Details	

Immunogen:	This ACBG2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 237-265 amino acids from the Central region of human ACBG2.
Clone:	RB24908
Isotype:	Ig Fraction
Purification:	This antibody is purified through a protein A column, followed by peptide affinity purification.

Target Details

Target:	ACSBG2
Alternative Name:	ACBG2 (ACSBG2 Products)

Target Details


Molecular Weight:	74354
Gene ID:	81616
NCBI Accession:	NP_112186
UniProt:	Q5FVE4
Pathways:	Monocarboxylic Acid Catabolic Process

Application Details

Application Notes:	WB: 1:1000. WB: 1:1000. IHC-P: 1:50~100. FC: 1:10~50
Restrictions:	For Research Use only


Handling

Format:	Liquid
Buffer:	Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Maintain refrigerated at 2-8 °C for up to 6 months. For long term storage store at -20 °C in small aliquots to prevent freeze-thaw cycles.
Expiry Date:	6 months




Flow Cytometry

Image 1. ACBG2 Antibody (Center) (ABIN653811 and ABIN2843086) flow cytometric analysis of K562 cells (right histogram) compared to a negative control cell (left histogram).FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.



Western Blotting

Image 2. Anti-ACBG2 Antibody (Center) at 1:1000 dilution + human testis lysate Lysates/proteins at 20 μg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 74 kDa Blocking/Dilution buffer: 5 % NFDM/TBST.



Western Blotting

Image 3. Western blot analysis of ACBG2 Antibody (Center) (ABIN653811 and ABIN2843086) in K562 cell line lysates (35 μ g/lane). ACBG2 (arrow) was detected using the purified Pab.

Please check the product details page for more images. Overall 4 images are available for ABIN653811.