antibodies - online.com







anti-B3GNT6 antibody (AA 146-173)

Images



\sim				
	$ V \cap$	r\/I	19	٨

Overview	
Quantity:	400 μL
Target:	B3GNT6
Binding Specificity:	AA 146-173
Reactivity:	Human, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This B3GNT6 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))
Product Details	
Immunogen:	This B3GNT6 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 146-173 amino acids from the Central region of human B3GNT6.
Clone:	RB24269
Isotype:	Ig Fraction
Purification:	This antibody is purified through a protein A column, followed by peptide affinity purification.
Target Details	
Target:	B3GNT6
Alternative Name:	B3GNT6 (B3GNT6 Products)
Molecular Weight:	42748

Target Details

Gene ID:	192134
NCBI Accession:	NP_619651
UniProt:	Q6ZMB0


Application Details

Application Notes:	WB: 1:1000. IHC-P: 1:50~100
Restrictions:	For Research Use only

Handling


Format:	Liquid
Buffer:	Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Maintain refrigerated at 2-8 °C for up to 6 months. For long term storage store at -20 °C in small aliquots to prevent freeze-thaw cycles.
Expiry Date:	6 months

Images



Immunohistochemistry (Paraffin-embedded Sections)

Image 1. B3GNT6 Antibody (Center) (ABIN653842 and ABIN2843106) immunohistochemistry analysis in formalin fixed and paraffin embedded human colon carcinoma followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the B3GNT6 Antibody (Center) for immunohistochemistry. Clinical relevance has not been evaluated.



Western Blotting

Image 2. Western blot analysis of B3GNT6 Antibody (Center) (ABIN653842 and ABIN2843106) in mouse brain tissue lysates (35 μ g/lane). B3GNT6 (arrow) was detected using the purified Pab.