



[Go to Product page](#)

Datasheet for ABIN653994

anti-CTDSPL antibody (N-Term)

2 Images

Overview

Quantity:	400 µL
Target:	CTDSPL
Binding Specificity:	AA 1-30, N-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CTDSPL antibody is un-conjugated
Application:	Flow Cytometry (FACS), Western Blotting (WB)

Product Details

Immunogen:	This CTDSPL antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 1-30 amino acids from the N-terminal region of human CTDSPL.
Clone:	RB27643
Isotype:	Ig Fraction
Purification:	This antibody is purified through a protein A column, followed by peptide affinity purification.

Target Details

Target:	CTDSPL
Alternative Name:	CTDSPL (CTDSPL Products)
Molecular Weight:	31129

Target Details

Gene ID:	10217
NCBI Accession:	NP_001008393 , NP_005799
UniProt:	O15194


Application Details

Application Notes:	WB: 1:1000. FC: 1:10~50
Restrictions:	For Research Use only

Handling


Format:	Liquid
Buffer:	Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C, -20 °C
Storage Comment:	Maintain refrigerated at 2-8 °C for up to 6 months. For long term storage store at -20 °C in small aliquots to prevent freeze-thaw cycles.
Expiry Date:	6 months

Images



Flow Cytometry

Image 1. CTDSPL Antibody (N-term) (ABIN653994 and ABIN2843931) flow cytometric analysis of HeLa cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.



Western Blotting

Image 2. All lanes : Anti-CTDSPL Antibody (N-term) at 1:1000 dilution Lane 1: Hela whole cell lysate Lane 2: K562 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 31 kDa Blocking/Dilution buffer: 5 % NFDN/TBST.