

Datasheet for ABIN654104  
**anti-TFB2M antibody (C-Term)**[Go to Product page](#)

2 Images

1 Publication

## Overview

Quantity:	400 µL
Target:	TFB2M
Binding Specificity:	AA 368-396, C-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This TFB2M antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

## Product Details

Immunogen:	This TFB2M antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 368-396 amino acids from the C-terminal region of human TFB2M.
Clone:	RB22408
Isotype:	Ig Fraction
Purification:	This antibody is purified through a protein A column, followed by peptide affinity purification.

## Target Details

Target:	TFB2M
Alternative Name:	TFB2M ( <a href="#">TFB2M Products</a> )
Molecular Weight:	45349

## Target Details

Gene ID:	64216
NCBI Accession:	<a href="#">NP_071761</a>
UniProt:	<a href="#">Q9H5Q4</a>

## Application Details


Application Notes:	WB: 1:2000. IHC-P: 1:50~100
Restrictions:	For Research Use only

## Handling

Format:	Liquid
Buffer:	Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Maintain refrigerated at 2-8 °C for up to 6 months. For long term storage store at -20 °C in small aliquots to prevent freeze-thaw cycles.
Expiry Date:	6 months


## Publications

Product cited in:	Alkhatabi, McLornan, Kulasekararaj, Malik, Seidl, Darling, Gaken, Mufti: "RPL27A is a target of miR-595 and may contribute to the myelodysplastic phenotype through ribosomal dysgenesis." in: <b>Oncotarget</b> , Vol. 7, Issue 30, pp. 47875-47890, (2018) ( <a href="#">PubMed</a> ).
-------------------	--



#### Immunohistochemistry (Paraffin-embedded Sections)

**Image 1.** TFB2M Antibody (C-term) (ABIN654104 and ABIN2843988) immunohistochemistry analysis in formalin fixed and paraffin embedded human heart carcinoma followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the TFB2M Antibody (C-term) for immunohistochemistry. Clinical relevance has not been evaluated.



#### Western Blotting

**Image 2.** All lanes : Anti-TFB2M Antibody (C-term) at 1:2000 dilution Lane 1: WiDr whole cell lysate Lane 2: HL-60 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 45 kDa Blocking/Dilution buffer: 5 % NFDm/TBST.