antibodies -online.com





anti-DIS3 antibody (AA 295-323)

2 Images



Go to	D		
	Pron	HCL	nane

Overview	
Quantity:	400 μL
Target:	DIS3
Binding Specificity:	AA 295-323
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This DIS3 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))
Product Details	
Immunogen:	This DIS3 antibody is generated from rabbits immunized with a KLH conjugated synthetic
	peptide between 295-323 amino acids from the Central region of human DIS3.
Clone:	RB24091
Isotype:	lg Fraction
Purification:	This antibody is purified through a protein A column, followed by peptide affinity purification.
Target Details	
Target:	DIS3
Alternative Name:	DIS3 (DIS3 Products)
Molecular Weight:	109003

Target Details

Gene ID:	22894
NCBI Accession:	NP_001121698, NP_055768
UniProt:	Q9Y2L1

UniProt:	Q9Y2L1
Application Details	
Application Notes:	WB: 1:1000. IHC-P: 1:50~100
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Format: Buffer:	Liquid Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.
Buffer:	Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.


aliquots to prevent freeze-thaw cycles.

Maintain refrigerated at 2-8 °C for up to 6 months. For long term storage store at -20 °C in small

Expiry Date: 6 months


Storage Comment:

Validation report #104401 for Cleavage Under Targets and Release Using Nuclease (CUT&RUN)



Immunohistochemistry (Paraffin-embedded Sections)

Image 1. DIS3 antibody (Center) (ABIN654155 and ABIN2844022) immunohistochemistry analysis in formalin fixed and paraffin embedded human testis carcinoma followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the DIS3 antibody (Center) for immunohistochemistry. Clinical relevance has not been evaluated.



Western Blotting

Image 2. All lanes: Anti-DIS3 Antibody (Center) at 1:1000 dilution Lane 1: Hela whole cell lysate Lane 2: Jurkat whole cell lysate Lysates/proteins at 20 μg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size: 109 kDa Blocking/Dilution buffer: 5 % NFDM/TBST.