

Datasheet for ABIN654195
anti-Radil antibody (C-Term)

2 Images

[Go to Product page](#)

Overview

Quantity:	400 µL
Target:	Radil
Binding Specificity:	AA 1044-1071, C-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Radil antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

Product Details

Immunogen:	This RADIL antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 1044-1071 amino acids from the C-terminal region of human RADIL.
Clone:	RB26913
Isotype:	Ig Fraction
Purification:	This antibody is purified through a protein A column, followed by peptide affinity purification.

Target Details

Target:	Radil
Alternative Name:	RADIL (Radil Products)
Molecular Weight:	117455

Target Details

Gene ID: 55698

NCBI Accession: [NP_060529](#)

UniProt: [Q96JH8](#)

Application Details

Application Notes: WB: 1:1000. IHC-P: 1:50~100

Restrictions: For Research Use only

Handling

Format: Liquid

Buffer: Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.

Preservative: Sodium azide


Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: 4 °C, -20 °C

Storage Comment: Maintain refrigerated at 2-8 °C for up to 6 months. For long term storage store at -20 °C in small aliquots to prevent freeze-thaw cycles.


Expiry Date: 6 months

Images



Immunohistochemistry (Paraffin-embedded Sections)

Image 1. RADIL antibody (C-term) (ABIN654195 and ABIN2844045) immunohistochemistry analysis in formalin fixed and paraffin embedded human pancreas tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the RADIL antibody (C-term) for immunohistochemistry. Clinical relevance has not been evaluated.



Western Blotting

Image 2. RADIL Antibody (C-term) (ABIN654195 and ABIN2844045) western blot analysis in SK-BR-3 cell line lysates (15 µg/lane). This demonstrates the RADIL antibody detected RADIL protein (arrow).