

Datasheet for ABIN654394
anti-Claudin 16 antibody (N-Term)



[Go to Product page](#)

3 Images

Overview

Quantity:	400 µL
Target:	Claudin 16 (CLDN16)
Binding Specificity:	AA 6-33, N-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Claudin 16 antibody is un-conjugated
Application:	Western Blotting (WB), Flow Cytometry (FACS), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

Product Details

Immunogen:	This CLDN16 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 6-33 amino acids from the N-terminal region of human CLDN16.
Clone:	RB30098
Isotype:	Ig Fraction
Purification:	This antibody is purified through a protein A column, followed by peptide affinity purification.

Target Details

Target:	Claudin 16 (CLDN16)
Alternative Name:	CLDN16 (CLDN16 Products)

Target Details

Background: Tight junctions represent one mode of cell-to-cell adhesion in epithelial or endothelial cell sheets, forming continuous seals around cells and serving as a physical barrier to prevent solutes and water from passing freely through the paracellular space. These junctions are comprised of sets of continuous networking strands in the outwardly facing cytoplasmic leaflet, with complementary grooves in the inwardly facing extracytoplasmic leaflet. The protein encoded by this gene, a member of the claudin family, is an integral membrane protein and a component of tight junction strands. It is found primarily in the kidneys, specifically in the thick ascending limb of Henle, where it acts as either an intercellular pore or ion concentration sensor to regulate the paracellular resorption of magnesium ions. Defects in this gene are a cause of primary hypomagnesemia, which is characterized by massive renal magnesium wasting with hypomagnesemia and hypercalciuria, resulting in nephrocalcinosis and renal failure. This gene and the CLDN1 gene are clustered on chromosome 3q28.

Molecular Weight: 33836

Gene ID: 10686

NCBI Accession: [NP_006571](#)

UniProt: [Q9Y517](#)

Pathways: [Hepatitis C](#)

Application Details

Application Notes: WB: 1:1000. IHC-P: 1:10~50. FC: 1:10~50

Restrictions: For Research Use only

Handling

Format: Liquid


Buffer: Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.

Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.


Storage: 4 °C, -20 °C

Storage Comment: Maintain refrigerated at 2-8 °C for up to 6 months. For long term storage store at -20 °C in small aliquots to prevent freeze-thaw cycles.




Immunohistochemistry (Paraffin-embedded Sections)

Image 1. CLDN16 antibody (N-term) (ABIN654394 and ABIN2844136) immunohistochemistry analysis in formalin fixed and paraffin embedded human Kidney tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the CLDN16 antibody (N-term) for immunohistochemistry. Clinical relevance has not been evaluated.



Western Blotting

Image 2. CLDN16 Antibody (N-term) (ABIN654394 and ABIN2844136) western blot analysis in MDA-M cell line lysates (35 µg/lane). This demonstrates the CLDN16 antibody detected the CLDN16 protein (arrow).



Flow Cytometry

Image 3. CLDN16 Antibody (N-term) (ABIN654394 and ABIN2844136) flow cytometric analysis of MDA-M cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.