

Datasheet for ABIN654462
anti-WDR48 antibody (C-Term)



[Go to Product page](#)

3 Images

Overview

Quantity:	400 µL
Target:	WDR48
Binding Specificity:	AA 603-630, C-Term
Reactivity:	Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This WDR48 antibody is un-conjugated
Application:	Western Blotting (WB), Flow Cytometry (FACS), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

Product Details

Immunogen:	This WDR48 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 603-630 amino acids from the C-terminal region of human WDR48.
Clone:	RB28145
Isotype:	Ig Fraction
Predicted Reactivity:	B, Pr
Purification:	This antibody is purified through a protein A column, followed by peptide affinity purification.

Target Details

Target:	WDR48
---------	-------

Target Details

Alternative Name: [WDR48 \(WDR48 Products\)](#)

Background: Regulator of deubiquitinating complexes. Acts as a strong activator of USP1 by enhancing the USP1-mediated deubiquitination of FANCD2, USP1 being almost inactive by itself. Also activates deubiquitinating activity of complexes containing USP12 and USP46, respectively. Activates deubiquitination by increasing the catalytic turnover without increasing the affinity of deubiquitinating enzymes for the substrate. In case of infection by Herpesvirus saimiri, may play a role in vesicular transport or membrane fusion events necessary for transport to lysosomes. Induces lysosomal vesicle formation via interaction with Herpesvirus saimiri tyrosine kinase-interacting protein (TIP). Subsequently, TIP recruits tyrosine-protein kinase LCK, resulting in down-regulation of T-cell antigen receptor TCR. May play a role in generation of enlarged endosomal vesicles via interaction with TIP. In case of infection by papillomavirus HPV11, promotes the maintenance of the viral genome via its interaction with HPV11 helicase E1.

Molecular Weight: 76210

Gene ID: 57599

NCBI Accession: [NP_065890](#)

UniProt: [Q8TAF3](#)

Application Details

Application Notes: WB: 1:1000. IHC-P: 1:50~100. FC: 1:10~50

Restrictions: For Research Use only

Handling

Format: Liquid



Buffer: Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.

Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: 4 °C,-20 °C

Storage Comment: Maintain refrigerated at 2-8 °C for up to 6 months. For long term storage store at -20 °C in small aliquots to prevent freeze-thaw cycles.



Flow Cytometry

Image 1. WDR48 Antibody (C-term) (ABIN654462 and ABIN2844195) flow cytometric analysis of 293 cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

Western Blotting

Image 2. Anti-WDR48_HUN at 1:1000 dilution + Rat testis lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 76 kDa Blocking/Dilution buffer: 5 % NFDM/TBST.

Immunohistochemistry (Paraffin-embedded Sections)

Image 3. WDR48 antibody (C-term) (ABIN654462 and ABIN2844195) immunohistochemistry analysis in formalin fixed and paraffin embedded human brain tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the WDR48 antibody (C-term) for immunohistochemistry. Clinical relevance has not been evaluated.