

Datasheet for ABIN654581
anti-PRRG3 antibody (AA 121-150)[Go to Product page](#)

2 Images

Overview

Quantity:	400 µL
Target:	PRRG3
Binding Specificity:	AA 121-150
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This PRRG3 antibody is un-conjugated
Application:	Western Blotting (WB), Flow Cytometry (FACS)

Product Details

Immunogen:	This PRRG3 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 121-150 amino acids from the Central region of human PRRG3.
Clone:	RB28384
Isotype:	Ig Fraction
Purification:	This antibody is purified through a protein A column, followed by peptide affinity purification.

Target Details

Target:	PRRG3
Alternative Name:	PRRG3 (PRRG3 Products)
Molecular Weight:	25875

Target Details

Gene ID: 79057

NCBI Accession: [NP_076987](#)

UniProt: [Q9BZD7](#)

Application Details

Application Notes: WB: 1:1000. FC: 1:10~50

Restrictions: For Research Use only

Handling

Format: Liquid

Buffer: Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.

Preservative: Sodium azide


Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: 4 °C,-20 °C

Storage Comment: Maintain refrigerated at 2-8 °C for up to 6 months. For long term storage store at -20 °C in small aliquots to prevent freeze-thaw cycles.


Expiry Date: 6 months

Images



Flow Cytometry

Image 1. PRRG3 Antibody (Center) (ABIN654581 and ABIN2844283) flow cytometric analysis of Jurkat cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.



Western Blotting

Image 2. Western blot analysis of lysates from human kidney and brain tissue lysates (from left to right), using PRRG3 Antibody (Center) (ABIN654581 and ABIN2844283). (ABIN654581 and ABIN2844283) was diluted at 1:1000 at each lane. A goat anti-rabbit IgG H&L(HRP) at 1:5000 dilution was used as the secondary antibody. Lysates at 35 µg per lane.