# antibodies - online.com







## anti-YdjC antibody (AA 169-198)



Molecular Weight:

34466



Overview	
Quantity:	400 μL
Target:	YdjC
Binding Specificity:	AA 169-198
Reactivity:	Human, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This YdjC antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))
Product Details	
Immunogen:	This YDJC antibody is generated from rabbits immunized with a KLH conjugated synthetic
	peptide between 169-198 amino acids from the Central region of human YDJC.
Clone:	RB28574
Isotype:	Ig Fraction
Purification:	This antibody is purified through a protein A column, followed by peptide affinity purification.
Target Details	
Target:	YdjC
Alternative Name:	YDJC (YdjC Products)

#### **Target Details**

Gene ID:	150223
NCBI Accession:	NP_001017964
UniProt:	A8MPS7


#### **Application Details**

Application Notes:	WB: 1:1000. IHC-P: 1:50~100
Restrictions:	For Research Use only

#### Handling

Format:	Liquid
Buffer:	Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Maintain refrigerated at 2-8 °C for up to 6 months. For long term storage store at -20 °C in small aliquots to prevent freeze-thaw cycles.
Expiry Date:	6 months

#### **Images**



### Immunohistochemistry (Paraffin-embedded Sections)

Image 1. YDJC Antibody (Center) (ABIN654597 and ABIN2844297) immunohistochemistry analysis in formalin fixed and paraffin embedded human tonsil tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the YDJC Antibody (Center) for immunohistochemistry. Clinical relevance has not been evaluated.

#### **Western Blotting**

**Image 2.** YDJC Antibody (Center) (ABIN654597 and ABIN2844297) western blot analysis in mouse testis tissue lysates (35  $\mu$ g/lane). This demonstrates the YDJC antibody detected the YDJC protein (arrow).