

Datasheet for ABIN654607

anti-HHLA2 antibody (AA 58-87)

7 Images

[Go to Product page](#)

Overview

Quantity:	400 µL
Target:	HHLA2
Binding Specificity:	AA 58-87
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This HHLA2 antibody is un-conjugated
Application:	Western Blotting (WB), Flow Cytometry (FACS), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

Product Details

Immunogen:	This HHLA2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 58-87 amino acids of human HHLA2.
Clone:	RB30079
Isotype:	IgG
Purification:	This antibody is purified through a protein A column, followed by peptide affinity purification.

Target Details

Target:	HHLA2
Alternative Name:	HHLA2 (HHLA2 Products)

Target Details

Molecular Weight:	46850
Gene ID:	11148
NCBI Accession:	NP_009003
UniProt:	Q9UM44


Application Details

Application Notes:	WB: 1:2000. WB: 1:2000. WB: 1:2000. WB: 1:2000. IHC-P: 1:50~100. IHC-P: 1:25. IHC-P: 1:25
Restrictions:	For Research Use only

Handling

Format:	Liquid
Buffer:	Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C, -20 °C
Storage Comment:	Maintain refrigerated at 2-8 °C for up to 6 months. For long term storage store at -20 °C in small aliquots to prevent freeze-thaw cycles.
Expiry Date:	6 months

Images




Immunohistochemistry (Paraffin-embedded Sections)

Image 1. A staining HHLA2 in human pancreas tissue sections by Immunohistochemistry (IHC-P - paraformaldehyde-fixed, paraffin-embedded sections). Tissue was fixed with formaldehyde and blocked with 3 % BSA for 0.5 hour at room temperature, antigen retrieval was by heat mediation with a citrate buffer (pH 6). Samples were incubated with primary antibody (1/25) for 1 hour at 37 °C. A undiluted biotinylated goat polyvalent antibody was

used as the secondary antibody.


Western Blotting

Image 2. Anti-HHLA2 Antibody (N-term) at 1:2000 dilution + Human kidney tissue lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 70 kDa Blocking/Dilution buffer: 5 % NFDM/TBST.



Western Blotting

Image 3. All lanes : Anti-HHLA2 Antibody (N-term) at 1:2000 dilution Lane 1: Human kidney lysate Lane 2: A549 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 70 kDa Blocking/Dilution buffer: 5 % NFDM/TBST.



Please check the [product details page](#) for more images. Overall 7 images are available for ABIN654607.