



[Go to Product page](#)

Datasheet for ABIN654725

anti-CU070 Protein antibody (C-Term)

2 Images

Overview

Quantity:	400 µL
Target:	CU070 Protein (CU070)
Binding Specificity:	AA 146-174, C-Term
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CU070 Protein antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

Product Details

Immunogen:	This CU070 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 146-174 amino acids from the C-terminal region of human CU070.
Clone:	RB28654
Isotype:	Ig Fraction
Purification:	This antibody is purified through a protein A column, followed by peptide affinity purification.

Target Details

Target:	CU070 Protein (CU070)
Alternative Name:	CU070 (CU070 Products)
Molecular Weight:	25456

Target Details

Gene ID: 85395

NCBI Accession: [NP_478070](#)

UniProt: [Q9NSI2](#)

Application Details

Application Notes: WB: 1:1000. IHC-P: 1:50~100

Restrictions: For Research Use only

Handling

Format: Liquid

Buffer: Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.

Preservative: Sodium azide


Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: 4 °C,-20 °C

Storage Comment: Maintain refrigerated at 2-8 °C for up to 6 months. For long term storage store at -20 °C in small aliquots to prevent freeze-thaw cycles.


Expiry Date: 6 months

Images



Immunohistochemistry (Paraffin-embedded Sections)

Image 1. C antibody (C-term) (ABIN654725 and ABIN2844413) immunohistochemistry analysis in formalin fixed and paraffin embedded human skeletal muscle followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of C antibody (C-term) for immunohistochemistry. Clinical relevance has not been evaluated.



Western Blotting

Image 2. C Antibody (C-term) (ABIN654725 and ABIN2844413) western blot analysis in mouse stomach tissue lysates (35 µg/lane). This demonstrates the C antibody detected the C protein (arrow).