antibodies -online.com









Go to Product page

Overview	
Quantity:	400 μL
Target:	DNAJC22
Binding Specificity:	AA 247-276, C-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This DNAJC22 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))
Product Details	
Immunogen:	This DNAJC22 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 247-276 amino acids from the C-terminal region of human DNAJC22.
Clone:	RB27892
Isotype:	lg Fraction
Predicted Reactivity:	В
Purification:	This antibody is purified through a protein A column, followed by peptide affinity purification.
Target Details	
Target:	DNAJC22
Alternative Name:	DNAJC22 (DNAJC22 Products)

Target Details

Background:	May function as a co-chaperone.
Gene ID:	79962
NCBI Accession:	NP_079178
UniProt:	Q8N4W6

Application Details

Application Notes:	WB: 1:1000. IHC-P: 1:50~100
Restrictions:	For Research Use only

Handling


Format:	Liquid
Buffer:	Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Maintain refrigerated at 2-8 °C for up to 6 months. For long term storage store at -20 °C in small aliquots to prevent freeze-thaw cycles.
Expiry Date:	6 months

Images



Immunohistochemistry (Paraffin-embedded Sections)

Image 1. DNAJC22 antibody (C-term) (ABIN654825 and ABIN2844500) immunohistochemistry analysis in formalin fixed and paraffin embedded human kidney tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the DNAJC22 antibody (C-term) for immunohistochemistry. Clinical relevance has not been evaluated.



Western Blotting

Image 2. DNAJC22 Antibody (C-term) (ABIN654825 and ABIN2844500) western blot analysis in CEM cell line lysates (35 μ g/lane).This demonstrates the DNAJC22 antibody detected the DNAJC22 protein (arrow).