antibodies -online.com







anti-ZNF720 antibody (AA 34-62)

Images



Goto	Product	nage
00 10	iioaact	page

_					
U	V	er	VI	е	W

Overview	
Quantity:	400 μL
Target:	ZNF720
Binding Specificity:	AA 34-62
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This ZNF720 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))
Product Details	
Immunogen:	This ZNF720 antibody is generated from rabbits immunized with a KLH conjugated synthetic
	peptide between 34-62 amino acids from the Central region of human ZNF720.
Clone:	RB27926
Isotype:	lg Fraction
Purification:	This antibody is purified through a protein A column, followed by peptide affinity purification.
Target Details	
Target:	ZNF720
Alternative Name:	ZNF720 (ZNF720 Products)
Molecular Weight:	14454

Target Details

Gene ID:	124411
NCBI Accession:	NP_001124385
UniProt:	Q7Z2F6


Application Details

Application Notes:	WB: 1:1000. IHC-P: 1:50~100
Restrictions:	For Research Use only

Handling


Format:	Liquid
Buffer:	Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Maintain refrigerated at 2-8 °C for up to 6 months. For long term storage store at -20 °C in small aliquots to prevent freeze-thaw cycles.
Expiry Date:	6 months

Images



Immunohistochemistry (Paraffin-embedded Sections)

Image 1. ZN Antibody (Center) (ABIN654974 and ABIN2844614) immunohistochemistry analysis in formalin fixed and paraffin embedded human lung carcinoma followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the ZN Antibody (Center) for immunohistochemistry. Clinical relevance has not been evaluated.



Western Blotting

Image 2. ZN Antibody (Center) (ABIN654974 and ABIN2844614) western blot analysis in MCF-7 cell line lysates (35 μ g/lane).This demonstrates the ZN antibody detected the ZN protein (arrow).