

Datasheet for ABIN655008
anti-TEX37 antibody (AA 73-101)[Go to Product page](#)

2 Images

Overview

Quantity:	400 µL
Target:	TEX37
Binding Specificity:	AA 73-101
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This TEX37 antibody is un-conjugated
Application:	Western Blotting (WB), Flow Cytometry (FACS)

Product Details

Immunogen:	This C2orf51 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 73-101 amino acids from the Central region of human C2orf51.
Clone:	RB28739
Isotype:	Ig Fraction
Purification:	This antibody is purified through a protein A column, followed by peptide affinity purification.

Target Details

Target:	TEX37
Alternative Name:	C2orf51 (TEX37 Products)
Gene ID:	200523

Target Details

NCBI Accession: [NP_689883](#)

UniProt: [Q96LM6](#)

Application Details

Application Notes: WB: 1:1000. FC: 1:10~50

Restrictions: For Research Use only

Handling

Format: Liquid

Buffer: Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.

Preservative: Sodium azide


Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: 4 °C, -20 °C

Storage Comment: Maintain refrigerated at 2-8 °C for up to 6 months. For long term storage store at -20 °C in small aliquots to prevent freeze-thaw cycles.


Expiry Date: 6 months

Images



Flow Cytometry

Image 1. C2orf51 Antibody (Center) (ABIN655008 and ABIN2844643) flow cytometric analysis of NCI- cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.



Western Blotting

Image 2. All lanes : Anti-C2orf51 Antibody (Center) at 1:1000 dilution Lane 1: human testis lysate Lane 2: NCI- whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 21 kDa Blocking/Dilution buffer: 5 % NFDM/TBST.