

Datasheet for ABIN655063
anti-HPR antibody (AA 212-240)[Go to Product page](#)

4 Images

Overview

Quantity:	400 µL
Target:	HPR
Binding Specificity:	AA 212-240
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This HPR antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

Product Details

Immunogen:	This HPR antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 212-240 amino acids from the Central region of human HPR.
Clone:	RB18726
Isotype:	Ig Fraction
Purification:	This antibody is purified through a protein A column, followed by peptide affinity purification.

Target Details

Target:	HPR
Alternative Name:	HPR (HPR Products)
Background:	Ubiquitous transcription factor required for a diverse set of processes. It is a component of the

Target Details


	CCR4 complex involved in the control of gene expression (By similarity).
Molecular Weight:	39030
Gene ID:	3250
NCBI Accession:	NP_066275
UniProt:	P00739
Pathways:	Response to Water Deprivation

Application Details

Application Notes:	WB: 1:2000. WB: 1:8000. WB: 1:1000. IHC-P: 1:50~100
Restrictions:	For Research Use only


Handling

Format:	Liquid
Buffer:	Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Maintain refrigerated at 2-8 °C for up to 6 months. For long term storage store at -20 °C in small aliquots to prevent freeze-thaw cycles.
Expiry Date:	6 months




Immunohistochemistry (Paraffin-embedded Sections)

Image 1. HPR Antibody (Center) (ABIN655063 and ABIN2844693) immunohistochemistry analysis in formalin fixed and paraffin embedded human hepatocarcinoma followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of HPR Antibody (Center) for immunohistochemistry. Clinical relevance has not been evaluated.



Western Blotting

Image 2. Anti-HPR Antibody (Center) at 1:2000 dilution + human plasma lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 39 kDa Blocking/Dilution buffer: 5 % NFDM/TBST.



Western Blotting

Image 3. Anti-HPR Antibody (Center) at 1:8000 dilution + human liver lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 39 kDa Blocking/Dilution buffer: 5 % NFDM/TBST.

Please check the [product details page](#) for more images. Overall 4 images are available for ABIN655063.