

Datasheet for ABIN655083
anti-INHBB antibody (C-Term)[Go to Product page](#)

4 Images

Overview

| | |
|----------------------|--|
| Quantity: | 400 µL |
| Target: | INHBB |
| Binding Specificity: | AA 369-397, C-Term |
| Reactivity: | Human, Mouse |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This INHBB antibody is un-conjugated |
| Application: | Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)) |

Product Details

| | |
|-----------------------|--|
| Immunogen: | This INHBB antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 369-397 amino acids from the C-terminal region of human INHBB. |
| Clone: | RB23814 |
| Isotype: | Ig Fraction |
| Predicted Reactivity: | B, C, Pig, Rat |
| Purification: | This antibody is purified through a protein A column, followed by peptide affinity purification. |

Target Details

| | |
|-------------------|--|
| Target: | INHBB |
| Alternative Name: | INHBB (INHBB Products) |

Target Details

Background: The inhibin beta B subunit joins the alpha subunit to form a pituitary FSH secretion inhibitor. Inhibin has been shown to regulate gonadal stromal cell proliferation negatively and to have tumour-suppressor activity. In addition, serum levels of inhibin have been shown to reflect the size of granulosa-cell tumors and can therefore be used as a marker for primary as well as recurrent disease. Because expression in gonadal and various extragonadal tissues may vary severalfold in a tissue-specific fashion, it is proposed that inhibin may be both a growth/differentiation factor and a hormone. Furthermore, the beta B subunit forms a homodimer, activin B, and also joins with the beta A subunit to form a heterodimer, activin AB, both of which stimulate FSH secretion.

Molecular Weight: 45122

Gene ID: 3625

NCBI Accession: [NP_002184](#)

UniProt: [P09529](#)

Pathways: [Peptide Hormone Metabolism](#), [Hormone Activity](#), [Negative Regulation of Hormone Secretion](#)

Application Details

Application Notes: WB: 1:4000. WB: 1:1000. WB: 1:2000. IHC-P: 1:10~50

Restrictions: For Research Use only

Handling

Format: Liquid

Buffer: Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.


Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: 4 °C,-20 °C


Storage Comment: Maintain refrigerated at 2-8 °C for up to 6 months. For long term storage store at -20 °C in small aliquots to prevent freeze-thaw cycles.

Expiry Date: 6 months




Immunohistochemistry (Paraffin-embedded Sections)

Image 1. INHBB Antibody (C-term) (ABIN655083 and ABIN2844717) immunohistochemistry analysis in formalin fixed and paraffin embedded human liver tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of INHBB Antibody (C-term) for immunohistochemistry. Clinical relevance has not been evaluated.



Western Blotting

Image 2. All lanes : Anti-INHBB Antibody (C-term at 1:4000 dilution Lane 1: A549 whole cell lysate Lane 2: PC-3 whole cell lysate Lane 3: Mouse testis lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 45 kDa Blocking/Dilution buffer: 5 % NFDm/TBST.



Western Blotting

Image 3. All lanes : Anti-INHBB Antibody (C-term) at 1:1000 dilution Lane 1: A549 whole cell lysate Lane 2: SK-BR-3 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 45 kDa Blocking/Dilution buffer: 5 % NFDm/TBST.

Please check the [product details page](#) for more images. Overall 4 images are available for ABIN655083.