

Datasheet for ABIN655146
anti-GOLGA8A antibody (N-Term)

3 Images

[Go to Product page](#)

Overview

Quantity:	400 µL
Target:	GOLGA8A
Binding Specificity:	AA 129-157, N-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This GOLGA8A antibody is un-conjugated
Application:	Western Blotting (WB), Immunofluorescence (IF), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

Product Details

Immunogen:	This GOG8A antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 129-157 amino acids from the N-terminal region of human GOG8A.
Clone:	RB30142
Isotype:	Ig Fraction
Purification:	This antibody is purified through a protein A column, followed by peptide affinity purification.

Target Details

Target:	GOLGA8A
Alternative Name:	GOG8A (GOLGA8A Products)

Target Details

Molecular Weight:	70117
NCBI Accession:	NP_851422
UniProt:	A7E2F4


Application Details

Application Notes:	IF: 1:10~50. WB: 1:1000. IHC-P: 1:10~50
Restrictions:	For Research Use only

Handling



Format:	Liquid
Buffer:	Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Maintain refrigerated at 2-8 °C for up to 6 months. For long term storage store at -20 °C in small aliquots to prevent freeze-thaw cycles.
Expiry Date:	6 months

Images



Immunohistochemistry (Paraffin-embedded Sections)

Image 1. GOG8A Antibody (N-term) (ABIN655146 and ABIN2844767) immunohistochemistry analysis in formalin fixed and paraffin embedded human liver tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of GOG8A Antibody (N-term) for immunohistochemistry. Clinical relevance has not been evaluated.



Immunofluorescence

Image 2. Confocal immunofluorescent analysis of GOG8A Antibody (N-term) (ABIN655146 and ABIN2844767) with 293 cell followed by Alexa Fluor 488-conjugated goat anti-rabbit IgG (green). DI was used to stain the cell nuclear (blue).

Western Blotting

Image 3. GOG8A Antibody (N-term) (ABIN655146 and ABIN2844767) western blot analysis in 293 cell line lysates (35 µg/lane). This demonstrates the GOG8A antibody detected the GOG8A protein (arrow).