

Datasheet for ABIN655197

**anti-FAM96A antibody (C-Term)**

3 Images

[Go to Product page](#)

## Overview

Quantity:	400 µL
Target:	FAM96A
Binding Specificity:	AA 109-138, C-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This FAM96A antibody is un-conjugated
Application:	Western Blotting (WB), Immunofluorescence (IF), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

## Product Details

Immunogen:	This FA96A antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 109-138 amino acids from the C-terminal region of human FA96A.
Clone:	RB28646
Isotype:	Ig Fraction
Predicted Reactivity:	B, M
Purification:	This antibody is purified through a protein A column, followed by peptide affinity purification.

## Target Details

Target:	FAM96A
---------	--------

## Target Details


Alternative Name:	FA96A ( <a href="#">FAM96A Products</a> )
Molecular Weight:	18355
Gene ID:	84191
NCBI Accession:	<a href="#">NP_001014812</a> , <a href="#">NP_115607</a>
UniProt:	<a href="#">Q9H5X1</a>

## Application Details

Application Notes:	IF: 1:10~50. WB: 1:2000. IHC-P: 1:50~100
Restrictions:	For Research Use only


## Handling

Format:	Liquid
Buffer:	Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Maintain refrigerated at 2-8 °C for up to 6 months. For long term storage store at -20 °C in small aliquots to prevent freeze-thaw cycles.
Expiry Date:	6 months




### Immunohistochemistry (Paraffin-embedded Sections)

**Image 1.** FA96A Antibody (C-term) (ABIN655197 and ABIN2844813) immunohistochemistry analysis in formalin fixed and paraffin embedded human kidney tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of FA96A Antibody (C-term) for immunohistochemistry. Clinical relevance has not been evaluated.



### Immunofluorescence

**Image 2.** Confocal immunofluorescent analysis of FA96A Antibody (C-term) (ABIN655197 and ABIN2844813) with WiDr cell followed by Alexa Fluor 488-conjugated goat anti-rabbit IgG (green). DAPI was used to stain the cell nuclear (blue).



### Western Blotting

**Image 3.** Anti-FA96A Antibody (C-term) at 1:2000 dilution + K562 whole cell lysate. Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 18 kDa. Blocking/Dilution buffer: 5 % NFDM/TBST.