antibodies - online.com







anti-SASH1 antibody (AA 596-625)

Images



Overview

Overview	
Quantity:	400 μL
Target:	SASH1
Binding Specificity:	AA 596-625
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This SASH1 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))
Product Details	
Immunogen:	This SASH1 antibody is generated from rabbits immunized with a KLH conjugated synthetic
	peptide between 596-625 amino acids from the Central region of human SASH1.
Clone:	RB18949
Isotype:	Ig Fraction
Purification:	This antibody is purified through a protein A column, followed by peptide affinity purification.
Target Details	
Target:	SASH1
Alternative Name:	SASH1 (SASH1 Products)
Background:	SASH1 may have a role in a signaling pathway and could act as a tumor suppressor.

Target Details

Molecular Weight:	136653
Gene ID:	23328
NCBI Accession:	NP_056093
UniProt:	O94885


Application Details

Application Notes:	IHC: 1:200. WB: 1:2000. IHC-P: 1:50~100
Restrictions:	For Research Use only

Handling


Format:	Liquid
Buffer:	Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Maintain refrigerated at 2-8 °C for up to 6 months. For long term storage store at -20 °C in small aliquots to prevent freeze-thaw cycles.
Expiry Date:	6 months

Images




Immunohistochemistry (Paraffin-embedded Sections)

Image 1. SASH1 Antibody (Center) (ABIN655510 and ABIN2845026) immunohistochemistry analysis in formalin fixed and paraffin embedded human placenta tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of SASH1 Antibody (Center) for immunohistochemistry. Clinical relevance has not been evaluated.



Immunohistochemistry

Image 2. Immunohistochemical analysis of paraffinembedded Human Colon cancer section using Pink1 . was diluted at 1:200 dilution. A undiluted biotinylated goat polyvalent antibody was used as the secondary, followed by DAB staining.



Western Blotting

Image 3. Anti-SASH1 Antibody (Center) at 1:2000 dilution + Mouse liver lysate Lysates/proteins at 20 μg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 137 kDa Blocking/Dilution buffer: 5 % NFDM/TBST.