



[Go to Product page](#)

Datasheet for ABIN655603
anti-AKR1C3 antibody (AA 107-135)

3 Images

Overview

| | |
|----------------------|--|
| Quantity: | 400 µL |
| Target: | AKR1C3 |
| Binding Specificity: | AA 107-135 |
| Reactivity: | Human |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This AKR1C3 antibody is un-conjugated |
| Application: | Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)) |

Product Details

| | |
|---------------|---|
| Immunogen: | This AKR1C3 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 107-135 amino acids from the Central region of human AKR1C3. |
| Clone: | RB31739 |
| Isotype: | Ig Fraction |
| Purification: | This antibody is purified through a protein A column, followed by peptide affinity purification. |

Target Details

| | |
|-------------------|---|
| Target: | AKR1C3 |
| Alternative Name: | AKR1C3 (AKR1C3 Products) |
| Background: | This gene encodes a member of the aldo/keto reductase superfamily, which consists of more |

Target Details

than 40 known enzymes and proteins. These enzymes catalyze the conversion of aldehydes and ketones to their corresponding alcohols by utilizing NADH and/or NADPH as cofactors. The enzymes display overlapping but distinct substrate specificity. This enzyme catalyzes the reduction of prostaglandin (PG) D2, PGH2 and phenanthrenequinone (PQ), and the oxidation of 9alpha,11beta-PGF2 to PGD2. It may play an important role in the pathogenesis of allergic diseases such as asthma, and may also have a role in controlling cell growth and/or differentiation. This gene shares high sequence identity with three other gene members and is clustered with those three genes at chromosome 10p15-p14.


| | |
|-------------------|---|
| Molecular Weight: | 36853 |
| Gene ID: | 8644 |
| NCBI Accession: | NP_001240837 , NP_003730 |
| UniProt: | P42330 |
| Pathways: | Retinoic Acid Receptor Signaling Pathway , Steroid Hormone Biosynthesis , Regulation of Hormone Metabolic Process , Regulation of Hormone Biosynthetic Process , C21-Steroid Hormone Metabolic Process , Protein targeting to Nucleus |

Application Details

| | |
|--------------------|--|
| Application Notes: | WB: 1:1000. WB: 1:1000. IHC-P: 1:10~50 |
| Restrictions: | For Research Use only |


Handling

| | |
|--------------------|--|
| Format: | Liquid |
| Buffer: | Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide. |
| Preservative: | Sodium azide |
| Precaution of Use: | This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only. |
| Storage: | 4 °C,-20 °C |
| Storage Comment: | Maintain refrigerated at 2-8 °C for up to 6 months. For long term storage store at -20 °C in small aliquots to prevent freeze-thaw cycles. |
| Expiry Date: | 6 months |




Immunohistochemistry (Paraffin-embedded Sections)

Image 1. AKR1C3 Antibody (Center) (ABIN655603 and ABIN2845087) immunohistochemistry analysis in formalin fixed and paraffin embedded human kidney tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of AKR1C3 Antibody (Center) for immunohistochemistry. Clinical relevance has not been evaluated.



Western Blotting

Image 2. AKR1C3 Antibody (Center) (ABIN655603 and ABIN2845087) western blot analysis in HeLa cell line lysates (35 µg/lane). This demonstrates the AKR1C3 antibody detected the AKR1C3 protein (arrow).



Western Blotting

Image 3. All lanes : Anti-AKR1C3 Antibody (Center) at 1:1000 dilution Lane 1: A549 whole cell lysate Lane 2: HepG2 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 37 kDa Blocking/Dilution buffer: 5 % NFDN/TBST.