



[Go to Product page](#)

Datasheet for ABIN655613  
**anti-PARP9 antibody (AA 599-629)**

2 Images

Overview

Quantity:	400 µL
Target:	PARP9
Binding Specificity:	AA 599-629
Reactivity:	Human, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This PARP9 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC)

Product Details

Immunogen:	This Parp9 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 599-629 amino acids of mouse Parp9.
Clone:	RB14182
Isotype:	IgG
Purification:	This antibody is purified through a protein A column, followed by peptide affinity purification.

Target Details

Target:	PARP9
Alternative Name:	Parp9 ( <a href="#">PARP9 Products</a> )
Background:	PARP9 is a novel risk related gene that is expressed at higher levels in fatal high risk diffuse

## Target Details

---

large B cell lymphomas.

Molecular Weight: 96659

Gene ID: 80285

NCBI Accession: [NP\\_084529](#)

UniProt: [Q8CAS9](#)

## Application Details

---

Application Notes: WB: 1:1000. IHC-P: 1:50~100

Restrictions: For Research Use only

## Handling

---

Format: Liquid

Buffer: Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.


Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: 4 °C,-20 °C


Storage Comment: Maintain refrigerated at 2-8 °C for up to 6 months. For long term storage store at -20 °C in small aliquots to prevent freeze-thaw cycles.

Expiry Date: 6 months



### Immunohistochemistry (Paraffin-embedded Sections)

**Image 1.** Parp9 Antibody (C-term) (ABIN655613 and ABIN2845094) immunohistochemistry analysis in formalin fixed and paraffin embedded human lymphoma followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of Parp9 Antibody (C-term) for immunohistochemistry. Clinical relevance has not been evaluated.



### Western Blotting

**Image 2.** All lanes : Anti-Parp9 Antibody (C-term) at 1:1000 dilution Lane 1: mouse lung lysate Lane 2: mouse thymus lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 97 kDa Blocking/Dilution buffer: 5 % NFDM/TBST.