



[Go to Product page](#)

Datasheet for ABIN655674
anti-GIN1 antibody (C-Term)

2 Images

Overview

Quantity:	400 µL
Target:	GIN1
Binding Specificity:	AA 464-492, C-Term
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This GIN1 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

Product Details

Immunogen:	This GIN1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 464-492 amino acids from the C-terminal region of human GIN1.
Clone:	RB29506
Isotype:	Ig Fraction
Predicted Reactivity:	Pr
Purification:	This antibody is purified through a protein A column, followed by peptide affinity purification.

Target Details

Target:	GIN1
Alternative Name:	GIN1 (GIN1 Products)

Target Details

Molecular Weight:	59842
Gene ID:	54826
NCBI Accession:	NP_060146
UniProt:	Q9NXP7


Application Details

Application Notes:	WB: 1:1000. IHC-P: 1:10~50
Restrictions:	For Research Use only

Handling


Format:	Liquid
Buffer:	Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Maintain refrigerated at 2-8 °C for up to 6 months. For long term storage store at -20 °C in small aliquots to prevent freeze-thaw cycles.
Expiry Date:	6 months

Images



Immunohistochemistry (Paraffin-embedded Sections)

Image 1. GIN1 Antibody (C-term) (ABIN655674 and ABIN2845140) immunohistochemistry analysis in formalin fixed and paraffin embedded human stomach tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of GIN1 Antibody (C-term) for immunohistochemistry. Clinical relevance has not been evaluated.



Western Blotting

Image 2. GIN1 Antibody (C-term) (ABIN655674 and ABIN2845140) western blot analysis in mouse stomach tissue lysates (35 µg/lane). This demonstrates the GIN1 antibody detected the GIN1 protein (arrow).