



[Go to Product page](#)

Datasheet for ABIN656377

## anti-RASSF10 antibody (AA 230-259)

2 Images

1 Publication

### Overview

Quantity:	400 µL
Target:	RASSF10
Binding Specificity:	AA 230-259
Reactivity:	Human, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This RASSF10 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

### Product Details

Immunogen:	This RASSF10 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 230-259 amino acids from the Central region of human RASSF10.
Clone:	RB31177
Isotype:	Ig Fraction
Purification:	This antibody is purified through a protein A column, followed by peptide affinity purification.

### Target Details

Target:	RASSF10
Alternative Name:	RASSF10 ( <a href="#">RASSF10 Products</a> )
Gene ID:	644943

## Target Details

---

NCBI Accession: [NP\\_001073990](#)

UniProt: [A6NK89](#)

## Application Details

---

Application Notes: WB: 1:1000. IHC-P: 1:10~50

Restrictions: For Research Use only

## Handling

---

Format: Liquid

Buffer: Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.

Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: 4 °C, -20 °C


Storage Comment: RASSF10 Antibody (Center) can be refrigerated at 2-8 °C for up to 6 months. For long term storage, place the at -20 °C.

Expiry Date: 6 months

## Publications


---

Product cited in: Mask, Schoenecker, Kane, Ransom, Bruemmer: "Serum antibody immunoreactivity to equine zona protein after SpayVac vaccination." in: **Theriogenology**, Vol. 84, Issue 2, pp. 261-7, (2015) ([PubMed](#)).



### Immunohistochemistry (Paraffin-embedded Sections)

**Image 1.** RASSF10 Antibody (Center) (ABIN656377 and ABIN2845672) immunohistochemistry analysis in formalin fixed and paraffin embedded human liver tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of RASSF10 Antibody (Center) for immunohistochemistry. Clinical relevance has not been evaluated.



### Western Blotting

**Image 2.** Anti-RASSF10 Antibody (Center) at 1:1000 dilution + HACAT whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 57 kDa Blocking/Dilution buffer: 5 % NFDm/TBST.