antibodies -online.com





anti-CHMP6 antibody (N-Term)



Image



Go to Product page

()	ve	K\ /		A .
	\cup	1 V/	Щ.	V۷

3.01.11311			
Quantity:	400 μL		
Target:	CHMP6		
Binding Specificity:	AA 1-30, N-Term		
Reactivity:	Human		
Host:	Rabbit		
Clonality:	Polyclonal		
Conjugate:	This CHMP6 antibody is un-conjugated		
Application:	Western Blotting (WB)		
Product Details			
Immunogen:	This CHMP6 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 1-30 amino acids from the N-terminal region of human CHMP6.		
Clone:	RB31342		
Isotype:	lg Fraction		
Purification:	This antibody is purified through a protein A column, followed by peptide affinity purification.		
Target Details			
Target:	CHMP6		
Alternative Name:	CHMP6 (CHMP6 Products)		
Background:	CHMP6 belongs to the chromatin-modifying protein/charged multivesicular body protein		

Target Details

(CHMP) family. These proteins are components of ESCRT-III (endosomal sorting complex required for transport III), a complex involved in degradation of surface receptor proteins and formation of endocytic multivesicular bodies (MVBs). Some CHMPs have both nuclear and cytoplasmic/vesicular distributions, and one such CHMP, CHMP1A (MIM 164010), is required for both MVB formation and regulation of cell cycle progression (Tsang et al., 2006 [PubMed 16730941]).

Molecular Weight: 23485

Gene ID: 79643

NCBI Accession: NP_078867

UniProt: Q96FZ7

Application Details

Application Notes: WB: 1:1000

storage, place the at -20 °C.

Restrictions: For Research Use only

Handling

Format:

Liquid

Buffer:

Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.

Preservative:

Sodium azide

Precaution of Use:

This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.


Storage:

4 °C,-20 °C

Storage Comment:

CHMP6 Antibody (N-term) can be refrigerated at 2-8 °C for up to 6 months. For long term

Expiry Date: 6 months



Western Blotting

Image 1. Western blot analysis of lysate from human spleen tissue lysate, using CH Antibody (N-term) (ABIN656387 and ABIN2845682). (ABIN656387 and ABIN2845682) was diluted at 1:1000 at each lane. A goat anti-rabbit IgG H&L(HRP) at 1:5000 dilution was used as the secondary antibody. Lysate at 35 μg per lane.