

Datasheet for ABIN656550
anti-MGAT4C antibody (C-Term)[Go to Product page](#)

2 Images

Overview

| | |
|----------------------|--|
| Quantity: | 400 µL |
| Target: | MGAT4C |
| Binding Specificity: | AA 303-332, C-Term |
| Reactivity: | Human |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This MGAT4C antibody is un-conjugated |
| Application: | Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)) |

Product Details

| | |
|-----------------------|--|
| Immunogen: | This MGAT4C antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 303-332 amino acids from the C-terminal region of human MGAT4C. |
| Clone: | RB32145 |
| Isotype: | Ig Fraction |
| Predicted Reactivity: | Pr, M, Pig |
| Purification: | This antibody is purified through a protein A column, followed by peptide affinity purification. |

Target Details

| | |
|-------------------|--|
| Target: | MGAT4C |
| Alternative Name: | MGAT4C (MGAT4C Products) |

Target Details

Background: Glycosyltransferase that participates in the transfer of N-acetylglucosamine (GlcNAc) to the core mannose residues of N-linked glycans. Catalyzes the formation of the GlcNAc β 1-4 branch on the GlcNAc β 1-2Man α 1-3 arm of the core structure of N-linked glycans. Essential for the production of tri- and tetra-antennary N-linked sugar chains (By similarity). Does not catalyze the transfer of GlcNAc to the Man α 1-6 arm to form GlcNAc β 1-4Man α 1-6 linkage ('GnT-VI' activity).

Molecular Weight: 56061

Gene ID: 25834

NCBI Accession: [NP_037376](#)

UniProt: [Q9UBM8](#)

Application Details

Application Notes: WB: 1:1000. IHC-P: 1:10~50

Restrictions: For Research Use only

Handling

Format: Liquid

Buffer: Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.


Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: 4 °C, -20 °C


Storage Comment: MGAT4C Antibody (C-term) can be refrigerated at 2-8 °C for up to 6 months. For long term storage, place the at -20 °C.

Expiry Date: 6 months



Immunohistochemistry (Paraffin-embedded Sections)

Image 1. MGAT4C Antibody (C-term) (ABIN656550 and ABIN2845812) immunohistochemistry analysis in formalin fixed and paraffin embedded human skeletal muscle followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of MGAT4C Antibody (C-term) for immunohistochemistry. Clinical relevance has not been evaluated.



Western Blotting

Image 2. MGAT4C Antibody (C-term) (ABIN656550 and ABIN2845812) western blot analysis in HepG2 cell line lysates (35 µg/lane). This demonstrates the MGAT4C antibody detected the MGAT4C protein (arrow).