



[Go to Product page](#)

Datasheet for ABIN656598

anti-Motilin Receptor antibody (C-Term)

2 Images

Overview

Quantity:	400 µL
Target:	Motilin Receptor (MLNR)
Binding Specificity:	AA 357-386, C-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Motilin Receptor antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

Product Details

Immunogen:	This MLNR antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 357-386 amino acids from the C-terminal region of human MLNR.
Clone:	RB31364
Isotype:	Ig Fraction
Purification:	This antibody is purified through a protein A column, followed by peptide affinity purification.

Target Details

Target:	Motilin Receptor (MLNR)
Alternative Name:	MLNR (MLNR Products)
Background:	MLNR is a receptor for motilin.

Target Details

Molecular Weight:	45344
Gene ID:	2862
NCBI Accession:	NP_001498
UniProt:	O43193


Application Details

Application Notes:	WB: 1:1000. IHC-P: 1:10~50
Restrictions:	For Research Use only

Handling


Format:	Liquid
Buffer:	Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	MLNR Antibody (C-term) can be refrigerated at 2-8 °C for up to 6 months. For long term storage, place the at -20 °C.
Expiry Date:	6 months

Images



Immunohistochemistry (Paraffin-embedded Sections)

Image 1. MLNR Antibody (C-term) (ABIN656598 and ABIN2845859) immunohistochemistry analysis in formalin fixed and paraffin embedded human stomach tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of MLNR Antibody (C-term) for immunohistochemistry. Clinical relevance has not been evaluated.



Western Blotting

Image 2. MLNR Antibody (C-term) (ABIN656598 and ABIN2845859) western blot analysis in NCI- cell line lysates (35 µg/lane). This demonstrates the MLNR antibody detected the MLNR protein (arrow).