antibodies -online.com







anti-TBL2 antibody (N-Term)





$C \circ + \circ$	Produc	+
(¬() (()	Promi	:i nane

Overview	
Quantity:	400 μL
Target:	TBL2
Binding Specificity:	AA 43-72, N-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This TBL2 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))
Product Details	
lmmunogen:	This TBL2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 43-72 amino acids from the N-terminal region of human TBL2.
Clone:	RB32541
Isotype:	Ig Fraction
Purification:	This antibody is purified through a protein A column, followed by peptide affinity purification.
Target Details	
Target:	TBL2
Alternative Name:	TBL2 (TBL2 Products)
Background:	This gene encodes a member of the beta-transducin protein family. Most proteins of the beta-

Target Details


	transducin family are involved in regulatory functions. This protein is possibly involved in some intracellular signaling pathway. This gene is deleted in Williams-Beuren syndrome, a developmental disorder caused by deletion of multiple genes at 7q11.23.
Molecular Weight:	49798
Gene ID:	26608
NCBI Accession:	NP_036585
UniProt:	Q9Y4P3


Application Details

Application Notes:	WB: 1:1000. IHC-P: 1:10~50
Restrictions:	For Research Use only

Handling

Format:	Liquid
Buffer:	Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	TBL2 Antibody (N-term) can be refrigerated at 2-8 °C for up to 6 months. For long term storage, place the at -20 °C.
Expiry Date:	6 months





Immunohistochemistry (Paraffin-embedded Sections)

Image 1. TBL2 Antibody (N-term) (ABIN656760 and ABIN2845981) immunohistochemistry analysis in formalin fixed and paraffin embedded human pancreas tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of TBL2 Antibody (N-term) for immunohistochemistry. Clinical relevance has not been evaluated.

Western Blotting

Image 2. TBL2 Antibody (N-term) (ABIN656760 and ABIN2845981) western blot analysis in ZR-75-1 cell line lysates (35 μ g/lane). This demonstrates the TBL2 antibody detected the TBL2 protein (arrow).