

Datasheet for ABIN656859

anti-Neuritin 1-Like antibody (AA 40-69)[Go to Product page](#)**2** Images

Overview

Quantity:	400 µL
Target:	Neuritin 1-Like (NRN1L)
Binding Specificity:	AA 40-69
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Neuritin 1-Like antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

Product Details

Immunogen:	This NRN1L antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 40-69 amino acids from the Central region of human NRN1L.
Clone:	RB28394
Isotype:	Ig Fraction
Purification:	This antibody is purified through a protein A column, followed by peptide affinity purification.

Target Details

Target:	Neuritin 1-Like (NRN1L)
Alternative Name:	NRN1L (NRN1L Products)
Molecular Weight:	17786

Target Details

Gene ID: 123904

NCBI Accession: [NP_940845](#)

UniProt: [Q496H8](#)

Application Details

Application Notes: WB: 1:1000. IHC-P: 1:10~50

Restrictions: For Research Use only

Handling

Format: Liquid

Buffer: Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.

Preservative: Sodium azide


Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: 4 °C,-20 °C

Storage Comment: NRN1L Antibody (Center) can be refrigerated at 2-8 °C for up to 6 months. For long term storage, place the at -20 °C.


Expiry Date: 6 months

Images



Immunohistochemistry (Paraffin-embedded Sections)

Image 1. NRN1L Antibody (Center) (ABIN656859 and ABIN2846064) immunohistochemistry analysis in formalin fixed and paraffin embedded human brain tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of NRN1L Antibody (Center) for immunohistochemistry. Clinical relevance has not been evaluated.



Western Blotting

Image 2. NRN1L Antibody (Center) (ABIN656859 and ABIN2846064) western blot analysis in MCF-7 cell line lysates (35 µg/lane). This demonstrates the NRN1L antibody detected the NRN1L protein (arrow).