



[Go to Product page](#)

Datasheet for ABIN656946
anti-DUX4 antibody (C-Term)

2 Images

Overview

Quantity:	400 µL
Target:	DUX4
Binding Specificity:	AA 376-404, C-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This DUX4 antibody is un-conjugated
Application:	Western Blotting (WB), Flow Cytometry (FACS)

Product Details

Immunogen:	This DUX4 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 376-404 amino acids from the C-terminal region of human DUX4.
Clone:	RB32790
Isotype:	Ig Fraction
Purification:	This antibody is purified through a protein A column, followed by peptide affinity purification.

Target Details

Target:	DUX4
Alternative Name:	DUX4 (DUX4 Products)
Background:	DUX4 may be involved in transcriptional regulation.

Target Details

Molecular Weight:	44940
Gene ID:	22947
NCBI Accession:	NP_001264985
UniProt:	Q9UBX2


Application Details

Application Notes:	WB: 1:2000. FC: 1:25
Restrictions:	For Research Use only

Handling

Format:	Liquid
Buffer:	Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C, -20 °C
Storage Comment:	DUX4 Antibody (C-term) can be refrigerated at 2-8 °C for up to 6 months. For long term storage, place the at -20 °C.
Expiry Date:	6 months


Images



Flow Cytometry

Image 1. Overlay histogram showing U-2 OS cells stained with (ABIN656946 and ABIN2846135)(green line). The cells were fixed with 2 % paraformaldehyde (10 min) and then permeabilized with 90 % methanol for 10 min. The cells were then incubated in 2 % bovine serum albumin to block non-specific protein-protein interactions followed by the antibody ((ABIN656946 and ABIN2846135), 1:25 dilution) for 60 min at 37 °C. The secondary antibody used was Goat-

Anti-Rabbit IgG, DyLight® 488 Conjugated Highly Cross-Adsorbed(1583138) at 1/200 dilution for 40 min at 37 °C. Isotype control antibody (blue line) was rabbit IgG1 (1 µg/1x10⁶ cells) used under the same conditions. Acquisition of >10,000 events was performed.



Western Blotting

Image 2. All lanes : Anti-DUX4 Antibody (C-term) at 1:2000 dilution Lane 1: A549 whole cell lysate Lane 2: HT-1080 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 45 kDa Blocking/Dilution buffer: 5 % NFDM/TBST.