antibodies -online.com





anti-MUC15 antibody (C-Term)

2 Images



Go to Product page

)\/(

Overview	
Quantity:	400 μL
Target:	MUC15
Binding Specificity:	AA 209-237, C-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))
Product Details	
Immunogen:	This MUC15 antibody is generated from rabbits immunized with a KLH conjugated synthetic
	peptide between 209-237 amino acids from the C-terminal region of human MUC15.
Clone:	RB32832
Isotype:	Ig Fraction
Purification:	This antibody is purified through a protein A column, followed by peptide affinity purification.
Target Details	
Target:	MUC15
Alternative Name:	MUC15 (MUC15 Products)
Background:	MUC15 may play a role in the cell adhesion to the extracellular matrix.
Molecular Weight:	36294

Target Details

Gene ID:	143662	
NCBI Accession:	NP_001128563, NP_001128564, NP_663625	
UniProt: Q8N387		


Application Details

Application Notes:	WB: 1:1000. IHC-P: 1:10~50
Restrictions:	For Research Use only

Handling


Format:	Liquid
Buffer:	Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	MUC15 Antibody (C-term) can be refrigerated at 2-8 °C for up to 6 months. For long term storage, place the at -20 °C.
Expiry Date:	6 months

Images



Immunohistochemistry (Paraffin-embedded Sections)

Image 1. MUC15 Antibody (C-term) (ABIN656956 and ABIN2846141) immunohistochemistry analysis in formalin fixed and paraffin embedded human liver tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of MUC15 Antibody (C-term) for immunohistochemistry. Clinical relevance has not been evaluated.



Western Blotting

Image 2. MUC15 Antibody (C-term) (ABIN656956 and ABIN2846141) western blot analysis in WiDr cell line lysates (35 μ g/lane).This demonstrates the MUC15 antibody detected the MUC15 protein (arrow).