



[Go to Product page](#)

Datasheet for ABIN656974
anti-SUSD2 antibody (C-Term)

2 Images

Overview

Quantity:	400 µL
Target:	SUSD2
Binding Specificity:	AA 716-745, C-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This SUSD2 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

Product Details

Immunogen:	This SUSD2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 716-745 amino acids from the C-terminal region of human SUSD2.
Clone:	RB32926
Isotype:	Ig Fraction
Purification:	This antibody is purified through a protein A column, followed by peptide affinity purification.

Target Details

Target:	SUSD2
Alternative Name:	SUSD2 (SUSD2 Products)
Molecular Weight:	90208

Target Details

Gene ID: 56241

NCBI Accession: [NP_062547](#)

UniProt: [Q9UGT4](#)

Application Details

Application Notes: WB: 1:1000. IHC-P: 1:10~50

Restrictions: For Research Use only

Handling

Format: Liquid

Buffer: Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.

Preservative: Sodium azide


Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: 4 °C, -20 °C

Storage Comment: SUSD2 Antibody (C-term) can be refrigerated at 2-8 °C for up to 6 months. For long term storage, place the at -20 °C.


Expiry Date: 6 months

Images



Immunohistochemistry (Paraffin-embedded Sections)

Image 1. SUSD2 Antibody (C-term) (ABIN656974 and ABIN2846158) immunohistochemistry analysis in formalin fixed and paraffin embedded human kidney tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of SUSD2 Antibody (C-term) for immunohistochemistry. Clinical relevance has not been evaluated.



Western Blotting

Image 2. Anti-SUSD2 Antibody (C-term) at 1:1000 dilution + WiDr whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 90 kDa Blocking/Dilution buffer: 5 % NFDM/TBST.