antibodies .- online.com





anti-GNAS antibody (C-Term)



Overview

Images



Go to Product page

Quantity:	400 μL
Target:	GNAS
Binding Specificity:	AA 287-315, C-Term
Reactivity:	Human

Host:	Rabbit	
Clonality:	Polyclonal	

Conjugate:	This GNAS antibody is un-conjugated
Application:	Western Blotting (WB), Immunofluorescence (IF), Flow Cytometry (FACS),

Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

Product Details

Immunogen:	This GNAS antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 287-315 amino acids from the C-terminal region of human GNAS.	
Clone:	RB33001	
Isotype:	IgG	
Predicted Reactivity:	M, Rat, B, Ha, Pig	
Purification:	This antibody is purified through a protein A column, followed by peptide affinity purification.	

Target Details

Target: GNAS

Target Details


Alternative Name:	GNAS (GNAS Products)	
Background:	Guanine nucleotide-binding proteins (G proteins) are involved as modulators or transducers in	
	various transmembrane signaling systems. The Gs protein is involved in hormonal regulation of	
	adenylate cyclase: it activates the cyclase in response to beta-adrenergic stimuli. Alternative	
	splicing of downstream exons of the GNAS gene is observed, which results in different forms of	
	the stimulatory G protein alpha subunit, a key element of the classical signal transduction	
	pathway linking receptor-ligand interactions with the activation of adenylyl cyclase and a variety	
	of cellular reponses. Multiple transcript variants have been found for this gene, but the full-	
	length nature and/or biological validity of some variants have not been determined. Mutations	
	in this gene result in pseudohypoparathyroidism type 1a, pseudohypoparathyroidism type 1b,	
	Albright hereditary osteodystrophy, pseudopseudohypoparathyroidism, McCune-Albright	
	syndrome, progressive osseus heteroplasia, polyostotic fibrous dysplasia of bone, and some pituitary tumors.	
Molecular Weight:	44250	
Gene ID:	2778	
UniProt:	Q5FWY2	
Pathways:	Thyroid Hormone Synthesis, cAMP Metabolic Process, Myometrial Relaxation and Contraction,	
	Embryonic Body Morphogenesis	
Application Details		
Application Notes:	IF: 1:10~50. WB: 1:1000. WB: 1:1000. IHC-P: 1:10~50. FC: 1:10~50	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Buffer:	Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.	
Preservative:	Sodium azide	
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which	
	should be handled by trained staff only.	
Storage:	4 °C,-20 °C	
Storage Comment:	GNAS Antibody (C-term) can be refrigerated at 2-8 °C for up to 6 months. For long term storage	

place the at -20 °C.

Expiry Date:


6 months

Images




Immunohistochemistry (Paraffin-embedded Sections)

Image 1. GNAS Antibody (C-term) (ABIN656989 and ABIN2850497) immunohistochemistry analysis in formalin fixed and paraffin embedded human pancreas tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of GNAS Antibody (C-term) for immunohistochemistry. Clinical relevance has not been evaluated.



Western Blotting

Image 2. Western blot analysis of GNAS (arrow) using rabbit polyclonal GNAS Antibody (C-term) (ABIN656989 and ABIN2850497). 293 cell lysates (2 μg/lane) either nontransfected (Lane 1) or transiently transfected (Lane 2) with the GNAS gene.



Western Blotting

Image 3. GNAS Antibody (C-term) (ABIN656989 and ABIN2850497) western blot analysis in 293,NCI- cell line lysates (35 μ g/lane).This demonstrates the GNAS antibody detected the GNAS protein (arrow).

Please check the product details page for more images. Overall 5 images are available for ABIN656989.