



[Go to Product page](#)

Datasheet for ABIN657091
anti-CNTLN antibody (N-Term)

2 Images

Overview

Quantity:	400 µL
Target:	CNTLN
Binding Specificity:	AA 199-227, N-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

Product Details

Immunogen:	This CNTLN antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 199-227 amino acids from the N-terminal region of human CNTLN.
Clone:	RB32726
Isotype:	Ig Fraction
Purification:	This antibody is purified through a protein A column, followed by peptide affinity purification.

Target Details

Target:	CNTLN
Alternative Name:	CNTLN (CNTLN Products)
Background:	The specific function of this protein remains unknown.
Molecular Weight:	161571

Target Details

NCBI Accession: [NP_001107867](#), [NP_060208](#)

UniProt: [Q9NXG0](#)

Application Details

Application Notes: WB: 1:1000. IHC-P: 1:10~50

Restrictions: For Research Use only

Handling

Format: Liquid

Buffer: Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.


Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: 4 °C, -20 °C


Expiry Date: 6 months

Images



Immunohistochemistry (Paraffin-embedded Sections)

Image 1. CNTLN Antibody (N-term) (ABIN657091 and ABIN2846249) immunohistochemistry analysis in formalin fixed and paraffin embedded human liver tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of CNTLN Antibody (N-term) for immunohistochemistry. Clinical relevance has not been evaluated.



Western Blotting

Image 2. Anti-CNTLN Antibody (N-term) at 1:1000 dilution + HL-60 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 162 kDa Blocking/Dilution buffer: 5 % NFDm/TBST.