



[Go to Product page](#)

Datasheet for ABIN657286  
**anti-CCDC120 antibody (N-Term)**

2 Images

### Overview

Quantity:	400 µL
Target:	CCDC120
Binding Specificity:	AA 59-88, N-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Application:	Western Blotting (WB)

### Product Details

Immunogen:	This CCDC120 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 59-88 amino acids from the N-terminal region of human CCDC120.
Clone:	RB33566
Isotype:	Ig Fraction
Purification:	This antibody is purified through a protein A column, followed by peptide affinity purification.

### Target Details

Target:	CCDC120
Alternative Name:	CCDC120 ( <a href="#">CCDC120 Products</a> )
Background:	This gene encodes a protein that contains a coiled-coil domain. Four alternatively spliced transcript variants that encode different protein isoforms have been described for this gene.

## Target Details

Molecular Weight:	67567
NCBI Accession:	<a href="#">NP_001156793</a> , <a href="#">NP_001156794</a> , <a href="#">NP_001156795</a> , <a href="#">NP_001258764</a> , <a href="#">NP_001258765</a> , <a href="#">NP_296375</a>
UniProt:	<a href="#">Q96HB5</a>


## Application Details

Application Notes:	WB: 1:1000. WB: 1:1000
Restrictions:	For Research Use only

## Handling


Format:	Liquid
Buffer:	Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Expiry Date:	6 months

## Images



### Western Blotting

**Image 1.** CCD Antibody (N-term) (ABIN657286 and ABIN2846373) western blot analysis in uterus tumor cell line lysates (35 µg/lane). This demonstrates the CCD antibody detected the CCD protein (arrow).



### Western Blotting

**Image 2.** All lanes : Anti-CCD Antibody (N-term) at 1:1000 dilution Lane 1: A431 whole cell lysate Lane 2: HeLa whole cell lysate Lysates/proteins at 20  $\mu$ g per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 68 kDa Blocking/Dilution buffer: 5 % NFDN/TBST.