

Datasheet for ABIN657379
anti-TXNDC15 antibody (C-Term)[Go to Product page](#)

2 Images

Overview

Quantity:	400 µL
Target:	TXNDC15
Binding Specificity:	AA 257-285, C-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This TXNDC15 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

Product Details

Immunogen:	This TXNDC15 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 257-285 amino acids from the C-terminal region of human TXNDC15.
Clone:	RB32916
Isotype:	Ig Fraction
Predicted Reactivity:	M
Purification:	This antibody is purified through a protein A column, followed by peptide affinity purification.

Target Details

Target:	TXNDC15
Alternative Name:	TXNDC15 (TXNDC15 Products)

Target Details



Background:	The specific function of the protein remains unknown.
Molecular Weight:	39885
Gene ID:	79770
NCBI Accession:	NP_078991
UniProt:	Q96J42

Application Details

Application Notes:	WB: 1:1000. IHC-P: 1:10~50
Restrictions:	For Research Use only

Handling

Format:	Liquid
Buffer:	Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	TXNDC15 Antibody (C-term) can be refrigerated at 2-8 °C for up to 6 months. For long term storage, place the at -20 °C.
Expiry Date:	6 months



Immunohistochemistry (Paraffin-embedded Sections)

Image 1. TXNDC15 Antibody (C-term) (ABIN657379 and ABIN2846424) immunohistochemistry analysis in formalin fixed and paraffin embedded human kidney tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of TXNDC15 Antibody (C-term) for immunohistochemistry. Clinical relevance has not been evaluated.

Western Blotting

Image 2. TXNDC15 Antibody (C-term) (ABIN657379 and ABIN2846424) western blot analysis in MCF-7 cell line lysates (35 µg/lane). This demonstrates the TXNDC15 antibody detected the TXNDC15 protein (arrow).