antibodies - online.com







anti-TRIM39 antibody (AA 248-277)





Overview	
Quantity:	400 μL
Target:	TRIM39
Binding Specificity:	AA 248-277
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This TRIM39 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))
Product Details	
Immunogen:	This TRIM39 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 248-277 amino acids from the Central region of human TRIM39.
Clone:	RB33252
Isotype:	lg Fraction
Purification:	This antibody is purified through a protein A column, followed by peptide affinity purification.
Target Details	
Target:	TRIM39
Alternative Name:	TRIM39 (TRIM39 Products)
Background:	The protein encoded by this gene is a member of the tripartite motif (TRIM) family. The TRIM

Target Details

	motif includes three zinc-binding domains, a RING, a B-box type 1 and a B-box type 2, and a
	coiled-coil region. The function of this protein has not been identified. This gene lies within the
	major histocompatibility complex class I region on chromosome 6. Alternate splicing results in
	two transcript variants encoding different isoforms. [provided by RefSeq].
Molecular Weight:	59690
Gene ID:	56658
NCBI Accession:	NP_067076, NP_742013

Application Details


Q9HCM9


Application Notes:	WB: 1:1000. IHC-P: 1:10~50
Restrictions:	For Research Use only

Handling

UniProt:

Format:	Liquid
Buffer:	Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	TRIM39 Antibody (Center) can be refrigerated at 2-8 °C for up to 6 months. For long term storage, place the at -20 °C.
Expiry Date:	6 months





Immunohistochemistry (Paraffin-embedded Sections)

Image 1. TRIM39 Antibody (Center) (ABIN657389 and ABIN2846430) immunohistochemistry analysis in formalin fixed and paraffin embedded human testis tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of TRIM39 Antibody (Center) for immunohistochemistry. Clinical relevance has not been evaluated.

Western Blotting

Image 2. TRIM39 Antibody (Center) (ABIN657389 and ABIN2846430) western blot analysis in MDA-M cell line lysates (35 μ g/lane).This demonstrates the TRIM39 antibody detected the TRIM39 protein (arrow).